



Biostratigraphy and Palaeoenvironment of the Upper Jurassic-Lower Cretaceous of Antsalova-Morondava basin-Madagascar

Rasolofotiana Edmond^a, Ranaivosoa V.^B, Andriambelomanana M. C.^C, Randriamanana N. E.^d

^{a, b, c}Mention Sedimentary basins Evolution Conservation, Science and Technology Domain, University of Antananarivo BP 906, 101 Antananarivo, Madagascar, edmondra@yahoo.com, rnvoaj@gmail.com, miorachrisandria@gmail.com

^dMention Processes and Industrial Ecology, Science and Technology Domain, University of Antananarivo BP 906, 101 Antananarivo, Madagascar, andymyou@gmail.com

ABSTRACT

Biostratigraphy and palaeoenvironment of the Upper Jurassic - Lower Cretaceous in Antsalova site are determined by micropaleontological study. The sediments collected in the field will pass physico-chemical treatments and geochemical analyzes in the laboratory. The sedimentary deposits of the Antsalova site have delivered a microfaunistic association constituted by Foraminifera: 18 species of benthic Foraminifera, grouped into 6 Families and 2 species of Ostracoda belonging to a single Family. Planktonic Foraminifera are absent in all the formation. This formation is subdivided into three levels by these microfaunal associations encountered and corresponds to stratigraphic stages ranging from the Callovian to the Hauterivian. The major elements (SiO₂, CaO, MgO) of the sediments, as well as the microfauna encountered, indicate a shallow marine environment (0 to 200 meters), with normal salinity and a hot climate. Microfossils geographical distribution and resemblance observed in this site, in many deposits of Madagascar, and the different world countries are related to the Ocean opening and the Tethys subduction.

Keywords: Foraminifera-Ostracods-stratigraphy-geochemistry-paleogeography.

1. Introduction

The Jurassic - Cretaceous interval is an important period in biological diversification at which many geological phenomena have occurred. The sedimentary series relating to this period is widely represented in the northern part of the Morondava basin, particularly the Antsalova site. Research work on sedimentary geology and/or paleontology has been carried out in this basin, namely the geological synthesis and paleontological studies of sedimentary lands in Madagascar (Besairie and Collignon, 1966 - 1971), the biostratigraphy and sedimentology of the Callovian -Cenomanian in Antsalova region (Ranarison, 1988), the identification of Jurassic microfossils from the Morondava Basin (Andriamalala, 1995),... And in this area, microfossils remain important temporal and biostratigraphic markers in sedimentary soils. The objective of this work is to determine the stratigraphic limits for the Upper Jurassic-Lower Cretaceous sedimentary formation in the Antsalova site through the study of the faunal associations encountered. The principle consists an inventory of the microfossils (Foraminifera, Ostracods, etc.), studying faunal associations and carrying out a geochemical analysis of the sediments in order to provide details for the palaeoenvironment reconstitution, as well as the paleobiogeography in the site.

2. Geological context

The sedimentary series in the northern part of the Morondava basin, parallel extend the base following the NNW-SSE direction. The division of the basin into two North and South sub-basins is evidenced by the bundles of the Ilovo and Mandabe faults for the southern part and the Bekopaka Bemaraha fault in the North zone in the southern part of the basin, the contact of the series sedimentary with the basement is faulted, in a major direction NNE-SSW. Overall, this part has a relatively flat geomorphology, marked by coasts corresponding to the erosion or tectonic cliff facing the east (Razafimbelo, 1987).

* Corresponding author.:

E-mail address: edmondra@yahoo.com

The general morphology of this basin is related to its tectonic style. Antsalova forms a series of small hills in the Antsiny Depression and the Cretaceous Plateau. This formation is characterized by glauconious marls and Duvalia marls which are often very clayey.

3. Materials and methods

The study site is located 17 km south of Antsalova village, near Tsingy de Bemaraha National Parks, belonging to Antsalova District, Melaky region, Mahajanga Province. Samples were collected from the sediment in place, unaltered with geographic coordinates: 18 ° 41 '50.7" 'South latitude and 44 ° 37' 14.3 'East longitude.



Figure 1: Location of the study site (Data FTM, BD 500).

The sediments collected on the field will continue physicochemical treatments and geochemical analyzes in the laboratory. Sorting and counting of microfossils is necessary in order to establish the relative frequencies of the different taxa and also for statistical chi-square calculations (test of independence).

Independence test

The independence test makes it possible to verify the independence between the taxa (species) and the different levels (I, II and III) of the studied site. This test is intended to compare the value of X_c^2 calculated with X_c^2 threshold read in the chi-square table for a fixed risk of error and $(r_1-1) (r_2 -1)$ degrees of freedom.

Hypothesis

H_0 : taxa distributions and levels are two independent variables in the Antsalova site.

H_1 : Taxa distributions and levels are not independent in the Antsalova site.

Decision rule

The significance level is 0.01

If X_c^2 calculated is greater than X_c^2 threshold read in the chi-square table for a fixed risk of error and $(r_1-1) (r_2 -1)$ degrees of freedom, the null hypothesis is rejected.

If X_c^2 calculated is less than X_c^2 threshold read in the chi-square table for a fixed risk of error and $(r_1-1) (r_2 -1)$ degrees of freedom, the null hypothesis is accepted.

With: r_1 : site modality, r_2 : taxa modality by level.

Hypothesis

H_0 : taxa distributions and levels are two independent variables in the Antsalova site.

H_1 : Taxa distributions and levels are not independent in the Antsalova site.

4. Results and interpretation

Lithostratigraphy

The lithological section of Antsalova generally consists of greenish yellow marls with Belemnites and Ammonites, with a thickness of about 15.5 meters.

As part of this work, five samples were taken from this site: ATL 1, 2, 3, 4, 5. It is made up from bottom to top by:

ATL1: yellowish gray marl,

ATL2: yellow to green marl,

ATL3: greenish marl spotted with yellow,

ATL4: greenish friable marl with Belemnites and *Aptychus*,

ATL5: greenish Belemnite marl with grains of sand on the upper surface.






AGE	THICKNESS	SAMPLING NUMBER	PROFIL	LITHOLOGY
LOWER CRETACEOUS	3,30	ATL5		greenish marl to belemnites with sand grains on the upper surface
UPPER JURASSIC	6	ATL4		friable greenish marl to belemnites and <i>Aptychus</i>
	1,5	ATL3		friable marl greenish stained with yellow
CALLOVIAN	2,10	ATL2		yellow to green marl
	2,70	ATL1		grey to yellow marl

Figure 2: Lithological section of Antsalova (scale: 1/100, ATL: Antsalova)

Faunistic composition

The sedimentary deposits in the Antsalova site have yielded a microfaunal association consisting of Foraminifera: 18 species of benthic Foraminifera grouped together in 6 families and 2 species of Ostracods belonging to a single Family. Planktonic Foraminifera are absent throughout the formation.




Figure 2: The microfauna found in the Antsalova site

1. *Textularia* sp. (Textulariidae), 2. *Ammobaculites* sp. (Lituolidae), 3. *Astacolus mutilatus* (Nodosariidae), 4. *Astacolus* sp. (Nodosariidae), 5. *Lenticulina* sp. (Nodosariidae), 6. *Lenticulina subtilis* (Nodosariidae), 7. *Lenticulina fracta* (Nodosariidae), 8. *Lenticulina cf. secans var angulosa* (Nodosariidae), 9. *Lenticulina tsaramandrosoensis* (Nodosariidae), 10. *Lenticulina dosineontos* (Nodosariidae), 11. *Neoflabellina lanceolata* (Nodosariidae), 12. *Planularia madagascariensis* (Nodosariidae), 13. *Saracenaria tsaramandrosoensis* (Nodosariidae), 14. *Lingulina* sp. (Nodosariidae), 15. *Rectoglandulina brandi* (Glandilunidae), 16. *Vaginulina (Citharina) sparcostata* (Glandilunidae), 17. *Spirillina tenuissima* (Spirillinidae), 18. *Globulina* sp. (Polymorphinidae), 19. *Cytherella index* (Carapace, left view) (Cytherellidae), 20. *Cytherella ex gr. Pyriformis* (Carapace, left view) (Cytherellidae).

Other fossils

Macrofossils were found in the lithology of the site namely Cephalopods (*Hibolites semicanaliculatus*, *Actinocamax plenus*, *Duvalia lata*, *Megatheutis giganteus*) and Lamellibranchs (*Lamellaptychus lamellosus*)

Table 1: Distribution of Benthic Foraminifera and Ostracods in the Antsalova site.

Microfauna	Samples	ATL1	ATL2	ATL3	ATL4	ATL5
Benthic foraminifera hyaline		69.00	88.00	169.00	315.00	298.00
Benthic foraminifera agglutinated		0.00	0.00	0.00	0.00	59.00
Total number of Foraminifera		69.00	88.00	169.00	315.00	357.00
Ostracods		0.00	0.00	21.00	42.00	34.00

Samples taken from the Antsalova (ATL) site offer a great wealth of microfauna (Table 1). This site is marked by a gradual increase in the number of Foraminifera (69.00 to 357.00) going from the base of the formation (ATL1) to the top (ATL5). Benthic Foraminifera are very abundant throughout the formation studied; on the other hand, planktonic forms are absent. Four species of benthic Foraminifera are left in open nomenclature: *Textularia* sp., *Ammobaculites* sp., *Globulina* sp., *Lenticulina* sp. and Ostracods are very rare or even absent in some lower levels. The subdivision of the lithological section of Antsalova is determined from the faunal composition and the facies. From bottom to top, it is made up of three levels (I, II, III). The observed numbers (ni) and theoretical numbers (ni²) of microfauna species for each level are clearly very different (Table 2).

Table 2: Contingency table for the characteristic taxa of each level in the Antsalova site.

N°	SPECIES	I		II		III		TOTAL
		ni	ni'	ni	ni'	ni	ni'	
01	<i>Ammobaculites sp.</i>	0.00	11.00	0.00	10.00	33.00	11.00	33.00
02	<i>Textularia sp.</i>	0.00	9.00	0.00	8.00	26.00	9.00	26.00
03	<i>Astacolus mutilatis</i>	36.00	12.00	0.00	11.00	0.00	13.00	36.00
04	<i>Astacolus sp.</i>	0.00	13.00	0.00	12.00	39.00	13.00	38.00
05	<i>Lenticulina fracta</i>	0.00	18.00	52.00	16.00	0.00	18.00	52.00
06	<i>Lenticulina sp.</i>	91.00	64.00	96.00	57.00	0.00	65.00	187.00
07	<i>Lenticulina subtilis</i>	51.00	17.00	0.00	15.00	0.00	18.00	51.00
08	<i>Lenticulina dineontos</i>	0.00	8.00	0.00	7.00	23.00	8.00	23.00
09	<i>Lenticulina tsaramandroensis</i>	0.00	12.00	0.00	11.00	35.00	12.00	35.00
10	<i>Lenticulina secans var. angulosa</i>	0.00	11.00	0.00	10.00	31.00	11.00	31.00
11	<i>Planularia madagascariensis</i>	45.00	33.00	47.00	29.00	0.00	32.00	92.00
12	<i>Neoflabellina lanceolata</i>	42.00	14.00	0.00	13.00	0.00	15.00	42.00
13	<i>Globulina sp.</i>	61.00	44.00	41.00	40.00	27.00	45.00	129.00
14	<i>Saracenaria tsaramandroensis</i>	0.00	7.00	0.00	7.00	21.00	7.00	21.00
15	<i>Lingulina sp.</i>	0.00	12.00	36.00	11.00	0.00	13.00	36.00
16	<i>Vaginulina (Citharina) sparicostata</i>	0.00	9.00	0.00	8.00	26.00	9.00	26.00
17	<i>Rectoglandulina drandi</i>	0.00	7.00	0.00	6.00	19.00	7.00	19.00
18	<i>Spirillina tenuisima</i>	0.00	6.00	0.00	6.00	18.00	6.00	18.00
19	<i>Cytherella index</i>	21.00	15.00	24.00	14.00	0.00	16.00	45.00
20	<i>Cytherella ex gr. pyriformis</i>	0.00	25.00	18.00	23.00	55.00	25.00	73.00
TOTAL		347		314		353		1013

Geochemical analysis result

Analysis of major elements in samples from the Antsalova site formation (ATL) shows that the SiO₂ contents and CaO are very high, but the other elements are generally low (TiO₂, Al₂O₃, Fe₂O₃, MgO). (Table 3)

Table 3: Geochemical analysis results of samples

Samples(for each site)	Chemical elements in %							
	SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	MgO	CaO	Ignition loss	Humidity
ATL1	38.50	0.50	13.50	5.50	3.50	20.50	10.05	5.50
ATL2	37.05	0.30	14.50	6.50	3.05	22.30	12.50	5.05
ATL3	33.50	0.50	10.50	5.50	2.50	25.50	10.50	4.50
ATL4	35.05	0.30	3.05	5.05	2.05	30.50	9.05	0.50
ATL5	44.50	0.50	1.50	6.05	2.50	19.05	7.50	0.50

5. Discussions


Biostratigraphy

Biozonation based on benthic Foraminifera is closely facies dependent. It takes place at the regional level and in correspondence with macrofauna which also have a good power of biostratigraphic resolution. The faunistic associations encountered make it possible to subdivide the formation of the Antsalova site into three levels. These three levels correspond to stratigraphic stages ranging from the Callovian to the Hauterivian:

level I: Callovian,

level II: Oxfordian - Kimmeridgian - Portlandian (Upper Jurassic),

level III: Berriasien - Valanginien – Hauterivien (Lower Cretaceous). (Figure 3)



Call. : Callovien, Oxf. : Oxfordian, Kim. : Kimmeridgien, Bér. : Berriasien, Val. : Valanginien, Hau. : Hauterivien.

Figure 4: Stratigraphic distribution of the microfauna and macrofauna encountered.

The distribution of microfauna species in the three levels (I, II, III) of the Antsalova Formation is very different. Indeed, the species encountered in each level are not the same. In order to check the difference between the diversity and the distribution of species in these three levels, it is necessary to apply the "independence test"

H₀: Species distributions and levels are independent in the Antsalova formation.

H₁: Species distributions and levels are dependent in the Antsalova formation.

$$\chi^2_c(19) = \sum_{40}^1 \frac{(ni - ni')^2}{ni'} = 1202.04$$

The calculated value of χ^2_c is greater than the threshold χ^2_c value read from the chi-square table for a risk of error 0.01 and degree of freedom 38, the null hypothesis is rejected. The distribution of species and the different levels in the Antsalova site are very dependent. The variation of species in the three levels confirms the stratigraphic subdivision in the Antsalova Formation from the Callovian to the Hauterivian.

The stratigraphy of the sequence studied and the stratigraphy of Hourcq V. (1950) and M. Collignon (1959- 1960) in the Antsalova region show a great similarity of sediment formation. This formation is generally made up of marls: Bélemnites marl, Duvalia marl, Ammonites marl, Bélemnites marl and Aptychus marl. The variation is manifested in the thickness of the layers, particularly for the Upper Jurassic and Lower Cretaceous (Table 4).

Table 4: Stratigraphic comparison of the present study with the stratigraphy of Hourcq V. (1950) and Collignon M. (1959-1960)

Age	Level	ANTSALOVA	
		Present site	Hourcq V. 1950) et Collignon M. (1959- 1960)
Lower Cretaceous	III	Greenish marl with <i>Duvalia lata</i>	Marl with <i>Duvalia</i>
Upper Jurassic	II	Greenish marl with <i>Bélemnites</i> and <i>Aptychus</i>	Marl with infill from limestone bank to <i>Hibolites</i> and <i>Aptychus</i>
Callovian	I	Green to grey marl	Limestone marl <i>Macrocephalites</i>

The Upper Jurassic Formation of East Africa corresponds to much more clayey and marly facies (Rajaomazava, 1992). This Formation is compatible with the deposition environments of Madagascar drift from East Africa.

Paleoecology

The paleoecological study is determined by the fossil faunal association contained in the sediments. The variation of the deposition medium is characterized from the variations of the faunas.

Certain species can then be considered as witnesses of the sedimentation environment for facies fossils.

Level I

The Nodosariidae and Polymorphinidae association prefer relatively shallow areas of continental shelves.

The fairly well-preserved forms testify to low transport and low hydrodynamics.

The CaO content is quite low, which indicates a less carbonate rich environment and the increase in the MgO percentage confirms a less hot climate at this level.

Level II

A reduction in the number of species is marked in this level. In this case, the environment is less favorable to the development of microfauna. The paleoecological interpretation of the environment is based on data provided by the nature of the sediments. The Ostracod valves are fairly well preserved, which indicates low transport and low hydrodynamics. The CaO content increases, the environment is rich in carbonate, and the decrease in the MgO proportion indicates a hot climate.

Level III

This level is marked by the predominance of Foraminifera. The genus *Saracenaria* is characteristic of shallow water that is 110m deep.

Ostracod valves are fairly well preserved, indicating low transport and low hydrodynamics.

The CaO content decreases, the medium is less rich in carbonate and it is confirmed by the abundance of smooth forms. The quantity in MgO is low, it results in a hot climate.

Paleobiogeography

The geographic distribution of marine microfossils (Foraminifera, Ostracods) is related to the opening of the Ocean and the subduction of the Tethys. Indeed, one can see the similarity and evolution of organisms in each part after the dislocation of the continent.

Some Genres or species of microfossils encountered in this site have been found in numerous deposits of Madagascar and also described in different countries of the world (Table 5).

Table 5: Paleobiogeographic distribution of Foraminifera and Ostracodia.

Foraminiferes and Ostracods	Madagascar		Another countries (Espitalie et Sigal, N.Gekoff, 1963)
	Espitalie and Sigal, N.Grekoff, (1963)	Present site	
<i>Astacolus</i>	Malakialina, Port-Berger, Tsaramandroso,	Antsalova	Amérique
<i>Lenticulina</i>	Antorolava, Ambanjabe, Tsaramandroso, Port- Berger	Antsalova	Allemagne
<i>Saracenaria</i>	Tsaramandroso	Antsalova	Allemagne, Pays-Bas
<i>Planularia</i>	Malakialina	Antsalova	Angleterre
<i>Lingulina</i>	Malakialina	Antsalova	Allemagne, Pays-Bas, Texas
<i>Rectoglandulina</i>	Tsararivotra, Malakialina	Antsalova	Amérique du Nord
<i>Spirillina</i>	Mahajanga	Antsalova	Pologne
<i>Cytherella index</i>	Ihopy	Antsalova	Europe Occidentale, Jura Suisse
<i>Cytherella ex gr. pyriformis</i>	Andranomavo, Marobarria, Mahajanga, Tsararivotra	Antsalova	Bassin de Paris

Conclusion

The biostratigraphic study of the Antsalova fossil site allows us to determine 18 species of benthic Foraminifera and 2 species of Ostracods. Based on the stratigraphic distribution of faunal associations of microfossils and macrofossils discovered, the Antsalova marl formation is subdivided into three levels:

- level I: Callovian,

- level II: Oxfordian –Portlandian (Upper Jurassic),
- level III: Berriasien – Hauterivien (Lower Cretaceous).

The site is characterized by a shallow environment of 0 to 200m, hot with normal salinity. The geographic distribution of marine microfossils (Foraminifera, Ostracods) is related to the opening of the Ocean and the subduction of the Tethys. Microfossils are very perfectionist tools for determining the age and nature of the layers traversed by drilling during prospecting or oil exploration. They are also very interesting for the realization of major works: tunnels, dams, roads, ... and this to conclude that in terms of sedimentary rocks, remember that they have container (marl or limestone) and content (microfossils and / or macrofossils), object of this study, the marl according to the results of geochemical study shows a potential of direct valorization or transformed into materials and agricultural amendments. It should be noted that marl can be used as a crop amendment product in order to make the soils less acidic, hence their necessity in agriculture and especially for development in general.

REFERENCES

1. Andriamalala, F. 1995. Contribution to the Jurassic Microfossil Study of the Morondava Basin, DEA Thesis, Department of Paleontology, University of Antananarivo, Antananarivo.
2. Besairie, H. & Collignon, M. 1971. Geology of Madagascar. Sedimentary terrains. Annale de géologie de Madagascar, booklet XXXV.
3. Collignon, M. 1959. Atlas of the characteristic Fossils of Madagascar, Fascicle V.
4. Espitalie, J. & Sigal, J. 1963. Contribution to the study of Foraminifera from the Upper Jurassic and Neocomian of the Majunga basin. Annale de géologie de Madagascar, fascicle XXXII, 99 p., XXXVI pl.
5. Grekoff, N. 1963. Contribution to the study of Ostracods of the Middle Mesozoic (Bathonian-Valanginian) of the Majunga basin, Madagascar, Annale de géologie de Madagascar, fasc. XXXIII, 167 p., 10 pl.
6. Hourcq V. 1950. The sedimentary lands of the Morondava region. Annale de Géologie Service des Mines, Madagascar, 109 p.
7. Chamley H. 2000. Bases of sedimentology 2nd edition, Dunod, Paris 1987 420p
8. Loeblich A. & Tappan, H. 1987. Foraminiferal genera and their classification. Van Nostrand Reinhold, New York, vol. 1, 970 p.; flight. 2, 212 p., 847 pls.
9. Oertli H. J. 1971. Paleoecology of Ostracodes, vol.5, 953 p.
10. Razafimbelo E. 1987. The Morondava basin (Madagascar): Geological and structural synthesis, Doctoral thesis, University L. Pasteur, Strasbourg, 256 p.
11. Rajaomazava F. 1992. Study of the subsidence of the Morondava basin (Madagascar) within the framework of the geodynamic evolution of the East African margin, Center Géologique et Géophysique Université des Sciences et Techniques du Languedoc 34095 - Mont Pellicier, cedex 5 - France, 204 p.
12. Rasolofotiana E. 2016. Biostratigraphy and Comparative Paleoecology of the Upper Jurassic and Lower Cretaceous in the regions of Andranomavo (Mahajange basin) and Antsalova (Morondava basin), Madagascar, PhD thesis, University of Antananarivo, 189p, Antananarivo
13. Hutton J. 1726-1797. The "Principle of Actualism" Web page: <https://fr.wikipedia.org/wiki/James-Hutton>.
14. Beauchamp J., 2013. Coastal carbonate sedimentation.