



Intrauterine Drug Delivery System

*Jayashri .S. Kolsure^{*1} , Khalid .U.Shaikh² , Nandkishor.B. Bawage³, Shyamalila .B.Bawage.⁴*

¹. B- pharmacy final year Latur college of pharmacy ,Hasegaon,.

² Department of Pharmaceutics ,Latur college of pharmacy ,Hasegaon.

³. Department of pharmaceutical Analysis,Latur college of pharmacy ,Hasegaon.

⁴. Department of pharmacognosy, Latur college of pharmacy ,Hasegaon, Tq .Ausa,Dist.Latur -413512 , Maharashtra, India.

ABSTRACT

The unconventional drug shipping systems refers to the techniques method technologies.the unconventional drug transport structures for shipping of drug apart from conventional drug transport structures,the maximum vital traditional drug shipping systems is that intrauterine drug transport systems.the intrauterine gadgets is a small piece of plastic that is inserted via a clinician into the uterus to prevent pregnancy.there are numerous differing types, which can be powerful for three to ten years relying on type .the IUD is a protracted - appearing reversible approach of contraception.there are one-of-a-kind forms of IUD .one type consists of the hormone levonorgestrel and every other kind is copper IUD.the hormone is continuously launched into the uterus and the every other form of IUD which is T fashioned covered with copper.this T shaped IUD can be left in uterus for two-five years.An IUD is a superb options for women who want a tremendously effective,long time , without problems reversible method of contraception. it may be for women who cannot use positive hormonal methods like delivery manipulate capsules. those copper IUD used for emergency contraception. and those contraception is inserted within five days after unprotected intercourse ,it is extra powerful than taking emergency contraceptive pills. If an IUD is inserted inside 7 days after unprotected vaginal intercourse it can save you a being pregnant .An IUD is a small gentle T fashioned tool with the nylon string connected.And IUD is inserted into the uterus by means of a healthcare issuer or clinicians .the IUD may additionally purpose mild infection of the lining of the uterus in order that a fertilized egg can not implant. intervene with the movement of sperm or kill sperm pace of the movement of the egg thru the fallopian tube .the IUD is 98% powerful in stopping being pregnant.A clinicians need to insert an IUD .it is also achieved while you are in your length .the clinicians will perform a pelvic examination and take a look at to peer in which your uterus is positioned .they may then insert a speculum into your vagina to peer your cervix and then wash your cervix with an antiseptic solution .next an IUD is put into your uterus and the strings are cut just below your cervix. IUD insertion can be uncomfortable .you may need to take ache capsules earlier than you have got an IUD inserted .removal of an IUD need to also be performed by means of a clinicians.

Keywords :- Intra-uterine drug delivery systems, copper T etc.

INTRAUTERINE DRUG DELIVERY SYSTEM :-

INTRODUCTION :

Intrauterine device (IUD) is a small object this is inserted through the cervix and located within the uterus to prevent being pregnant. A small string hangs down from the IUD into the upper part of the vagina. The IUD is not major during sex. IUDs can last 1-10 years. They affect the actions of eggs and sperm to prevent fertilization. additionally they alternate the liner of the uterus and prevent implantation.IUDs are ninety nine.2-99.nine% powerful as birth manage. They do no longer defend in opposition to sexually transmitted infections, along with HIV/AIDS. Insertion of an IUD takes handiest approximately 5 to ten minutes. Your fitness care provider will first do a pelvic examination to measure the scale, form, and role of your uterus and different reproductive organs. Your fitness care issuer will then positioned antiseptic answer onto your cervix. The IUD might be inserted up via the outlet of your cervix into your uterus. it's far positioned interior using a unique applicator that continues the IUD flat and closed until it's miles at the top of your uterus. you may sense cramping, however it normally isn't a whole lot. After the IUD has been inserted, your fitness care company will cut the string on the cease of the IUD so that it is quick sufficient in which it might not hassle you or companion. it will be lengthy sufficient so that you can take a look at to make sure that the IUD is in location. Your fitness care company will then communicate with you approximately what you want to do about checking the string. An IUD prevents being pregnant by means of preventing sperm from attaining an egg that your ovaries have launched. It does this through not letting sperm go into the egg. An IUD also changes the lining of the uterus so an egg does not implant within the lining if it's been fertilized. consequently,

the egg has no area to develop. IUDs are the most effective shape of nonpermanent birth manage. they're more than ninety nine% effective. which means that if a hundred ladies use the copper IUD or the levonorgestrel IUD ,less than 1 woman turns into pregnant in a 12 months. each month a female's ovary produces an egg. whilst a person ejaculates, his semen contains millions of sperm. It takes only this type of sperm to fertilise an egg and begin a being pregnant. If a pair has intercourse (penis in vagina) and do now not take precautions, there is a high chance of pregnancy. using contraception lets in you to have intercourse and avoid an unintended pregnancy. ladies at low danger of contracting sexually transmitted infections girls who can't or do now not wash to take The pill. An intrauterine device (IUD) normally is a small, bendy plastic frame. It often has copper wire or copper sleeves on it. it is inserted into a female 欵機 vagina thru her uterus. nearly all manufacturers of IUDs have strings, or threads, tied to them. The strings hold through the outlet of the cervix into the vagina. A company can remove the IUD with the aid of pulling lightly at the strings with forceps.

HISTORY OF INTRAUTERINE GADGETS :-

consistent with famous legend, Arab buyers inserted small stones into the uteruses in their camels to prevent pregnancy throughout lengthy wasteland trips. The story turned into at the start a tall tale to entertain delegates at a scientific conference on circle of relatives making plans; although it was later repeated as truth, it has no recognised ancient foundation. Precursors to IUDs have been first advertised in 1902. advanced from stem pessaries (wherein the stem held the pessary in region over the cervix), the 'stem' on these devices definitely prolonged into the uterus itself. also referred to as interuterine devices (due to the fact they occupied each the vagina and the uterus), they'd excessive costs of contamination and had been condemned by means of the clinical community. the first intrauterine device (contained entirely in the uterus) turned into defined in a German booklet in 1909, despite the fact that the author seems to have never advertised his product. In 1929, Dr. Ernst Gr fenberg of Germany posted a document on an IUD made of silk suture. He had located a 3% pregnancy fee amongst 1,100 girls the usage of his ring. In 1930, Dr. Gr fenberg reported a lower being pregnant fee of one.6% among 600 ladies the use of an improved ring wrapped in silver cord. Unbeknownst to Dr. Gr fenberg, the silver twine become infected with 26% copper. Copper's function in increasing IUD efficacy might not be diagnosed until almost forty years later.In 1934, japanese doctor Tenrei Ota developed a version of the Gr fenberg ring that contained a supportive shape in the center. The addition of this imperative disc lowered the IUD's expulsion charge. these devices nonetheless had high prices of infection, and their use and improvement turned into similarly stifled by means of global warfare II politics: contraception was forbidden in each Nazi Germany and Axisallied Japan. The Western global did not examine of the paintings by means of Gr fenberg and Ota until nicely after the struggle ended.[6] the first plastic IUD, the Margulies Coil or Margulies Spiral, changed into brought in 1958. This device changed into fairly massive, inflicting pain to a huge percentage of ladies users, and had a hard plastic tail, causing pain to their male partners. The Lippes Loop, a barely smaller tool with a monofilament tail, become introduced in 1962 and won in popularity over the Margulies device.[5] The stainless steel single-ring IUD was advanced within the Seventies[7] and broadly utilized in China due to low manufacturing expenses. The chinese language government banned production of metal IUDs in 1993 due to high failure rates (up to ten% consistent with yr). Dr. Howard Tatum, inside the america, conceived the plastic T-fashioned IUD in 1968. shortly thereafter Dr. Jaime Zipper, in Chile, added the concept of including copper to the devices to enhance their contraceptive effectiveness.[5][10] It became determined that copper-containing devices will be made in smaller sizes with out compromising effectiveness, resulting in fewer facet outcomes including pain and bleeding. T-formed gadgets had lower prices of expulsion due to their greater similarity to the shape of the uterus. The poorly designed Dalkon protect plastic IUD (which had a multifilament tail) become synthetic by means of the A. H. Robins corporation and sold with the aid of Robins in the america for three and a half years from January 1971 via June 1974, earlier than sales have been suspended by way of Robins on June 28, 1974 at the request of the FDA due to safety issues following reviews of one hundred ten septic spontaneous abortions in women with the Dalkon guard in location, seven of whom had died. Robins stopped global sales of the Dalkon defend in April 1975. 2nd-era copper-T IUDs have been also added within the 1970s. these gadgets had higher surface areas of copper, and for the primary time consistently achieved effectiveness prices of extra than 99%. worldwide today, aside from the new GyneFix, this is the handiest sort of non-hormonal IUD to be had.

ANATOMY OF UTERUS :

The girl reproductive organs encompass the uterus ,fallopian tubes, and the ovaries . Their function,size,and the physiological adjustments of mensruation ,pregnancy,and menopause.the uterus is a secondary sex organ. Secondary sex organs are additives of the reproductive tract that mature at some point of puberty beneath the affect of intercourse hormones comprised of number one sex organs (the ovaries in ladies and the testes in males).they're chargeable for the upkeep and transportation of gamates .In this newsletter , we shall observe the composition and the medical significance of the uterus.

Its vasculature , lymphatic deliver and innervation may be included .

The uterus has three elements ;

- 1.Fundas: top of the uterus , above the access factor of the uterine tubes .
- 2.frame : ordinary web page for implantation of the blastocyst.
- 3.Cervix: decrease a part of uterus linking it with the vagina . this part is structurally and functionally specific to the uterus.

Function of Uterus :

The uterus performs an essential function in human replica .

It's miles the organ of wherein the implantation and nourishment of the Fertilized ovum takes place . moreover it enables pushing out the baby at some point of beginning thru muscle contraction.

It features to nourish and house a fertilised egg until the fetus, Or offsprings , is prepared to be introduced .

Advantages:

- highly effective in preventing pregnancy .
- less expensive .
- Does now not interrupt sex.
- Does now not require accomplice s involvement.
- May be used for a protracted period of time.
- May be used as an emergency technique of delivery manage.
- An IUD offers long -time period contraception for three to 5 yrs and is value powerful.
- while you are equipped to turn out to be pregnant,the IUD may be removed by means of a health care company.

Hazards:

1. high failure price .
2. Requires several days of abstinence .
3. Not suitable to ladies with medical contraindication to being pregnant.
4. Beside the point for women with very irregular menstruations.

Side outcomes of IUD :

- What are the feasible facet effects of use of the IUD ?
- with the copper IUD,menstrual pain and bleeding may additionally growth.
- Bleeding between intervals might also occurs.
- both effects are commonplace in the first few months of use.
- the quantity of menstrual bleeding and the period of the menstrual duration generally lower over time.
- Menstrual pain additionally usually decreases.
- ache and heavy bleeding commonly lower with 1 years of use.
- a few ladies's also may additionally have side effects associated with the hormone in the IUDs.
- these aspect effects may additionally encompass headaches, nausea, melancholy and breast tenderness.

ILLNESS OF I.U.D. :

- 1.ENDOMETRIOSIS
- 2.ADENOMYOSIS
- 3.FIBROIDS
- 4.ENDOMETRIAL HYPERPLASIA
- 5.ENDOMETRIAL

1. ENDOMETRIOSIS:

Sign and SYMPTOM of ENDOMETRIASIS :

1. Very painful menstrual cramps.
- 2.very heavy menstrual bleeding.
- 3.standard persistent pelvic ache .
- 4.Painful bowel actions.
- 5.painful intercourse.
- 6.ache after intercourse.
- 7.painful urination.
- 8.Diarrhoea and constipation .

.2.ADENOMYOSIS:

SIGN and SYMPTOM OF ADENOMYOSIS:

- 1.Prolonged ,heavy and clotted menstrual bleeding .
- 2.Severe cramps or knife -like pain during menstruation .
- 3.Abdominal pain .
- 4.Backache.
- 5.Painful intercourse .
- 6.Abdominal pressure.
- 7.Abdominal bloating .
- 8.Nausea and vomiting during period .

TERATMENT of ADENOMYOSIS:

Anti- inflammatory drugs :

Such as ibuprofen (Advil, Motrin IB, others), to control the pain. By starting an anti-inflammatory medicine one or two days before your periods begins and taking it during your period begins and it during your periods .you can reduce menstrual blood flow and relive pain.

Hormone Therapy :

Combined estrogen-progestin birth control pills or hormones containing patches .Progestin only contraception -the absence of your menstrual periods which might provide some relief.

Hysterectomy:

If your pain is severe and no other treatments have worked ,your doctor might surgery to remove your uterus . removing your ovaries is necessary to control adenomyosis.

3.FIBROIDS:

SIGN and SYMPTOM of FIBROIDS

- 1.Heavy monthly periods.
- 2.pain during sexual intercourse.
- 3.Pelvic pain and pressure.
- 4.Frequent need to urinate .
- 5.Constipation and Bloating.
- 6.Chewing ice.
- 7.Anemia

TREATMENT OF FIBROIDS:

1.HYSTERECTOMY:

- a. If the uterus is >10w size.
- b.Or signs that are due to the fibroids.
- c.fast increase.
- d.stomach and vaginal.

MYOMECTOMY:

- a.great for unmarried fibroids in a younger girls.
- b.50% come to hysterectomy inside 5 yrs.
- three.Hysteroscopic resection .
- Uterine artery embolization.(UAE)
- Clinical options
- GnRh analogue.
- Mirena.

4.ENDOMETRIAL HYPERPLASIA :

SIGN AND SYMPTOM of ENDOMETRIAL HYPERPLASIA:

- 1.Heavy bleeding at some point of the menstrual durations.
- 2.Menstrual bleeding lasting for extra than the same old 5 days.
- three.Shorter menstrual cycles,less than 21 days .
- 4.Bleeding even after menopause.
- 5.decrease belly ache.
- 6.ordinary temper swings.
- 7.intense cramping sort of ache.
- 8.hot flushes.

TREATMENT OF ENDOMETRIAL HYPERPLASIA:

In lots of cases, endometrial hyperplasia may be dealt with with progestin .Progestin is given orally ,in a shot , in an intrauterine device ,or as a vaginal cream. How a lot and the way long you take it depend upon your age and the kind of hyperplasia .remedy with progestin can also motive vaginal bleeding like a menstrual duration .when you have unusual hyperplasia , specifically complicated extraordinary hyperplasia , the threat of cancer is multiplied . Hysterectomy generally is the satisfactory remedy choice if you do no longer want to have any more youngsters.

5.ENDOMETRIAL most cancers:

Between 2 to 10 percentage of IUDs fall out, maximum commonly within the first 12 months of use. there's an expanded hazard of falling out if : -an IUD become inserted right after shipping.

- an IUD has fallen out formerly.
- you've got never been pregnant.

There may be an improved hazard of miscarriage if an IUD is left inside the uterus all through a pregnancy.

The chance of pelvic inflammatory disorder (PID) can be accelerated slightly in the course of the first month after an IUD inserted. After the primary month the risk of STI is associated with the exposure to STI, no longer the usage of an IUD.

An IUD, in uncommon cases, might also connect to or perforate the wall of the uterus. The IUD may additionally want to be eliminated through minor surgical operation.

With Nova T or Flexi T three hundred (copper) IUD you could experience greater bleeding and cramping with periods. every so often, mild bleeding and spotting can appear between durations.

With Mirena IUD (hormonal) you could in the beginning revel in changes in

bleeding sample, complications, breast tenderness, pimples, weight changes or temper modifications. those hormone adjustments commonly decrease over the years.

Conclusion of IUD :

The IUD is a superb form of contraception in case you are not vulnerable to getting a sexually transmitted disease. it's far quality for women who've already had youngsters and are in a regular courting with one companion. Many providers endorse the progestin IUD for women who suffer from extraordinarily heavy, extended, or painful menstruation as it tends to lighten their durations or maybe suppress them altogether. And due to the fact they lose less blood, ladies using this IUD are less in all likelihood to increase iron-deficiency anemia, a condition that can purpose fatigue and different symptoms. some research have located that ladies with copper IUDs tend to have a decrease danger of endometrial most cancers. And a few specialists suspect they may discover that the progestin IUD has the equal effect, when you consider that it really is the case for progestin-only contraceptives just like the minipill and the shot. An IUD helps prevent egg and sperm from meeting. A woman does not get pregnant if sperm does not meet the egg. An IUD may stop a fertilized egg from developing within the uterus. Nova T or Flexi T three hundred IUD is about 98.7% effective in stopping pregnancy. Irena IUD is 99 % powerful in stopping pregnancy. Your hazard for buying pregnant after the IUD has been inserted may be very low. however, there's usually a mild threat that you may get pregnant, because the IUD is not 100% effective. in case you do get pregnant while the IUD is in place, you want to get the IUD eliminated. You should also see your health care company to make sure that your being pregnant isn't outdoor of the uterus (ectopic being pregnant). if your being pregnant isn't ectopic, you can choose whether to continue the being pregnant. in case you pick out to retain it, your caregiver will put off the IUD if the strings are visible. she can eliminate the device effortlessly with out an invasive method. there may be a slight chance that removing the IUD will cause you to lose the pregnancy, however you're more likely to lose the baby to infection (and placed your own health at threat) if you maintain it in. inside the unlikely event that the IUD can't be effortlessly eliminated and also you choose to continue.

References :-

1. Birth Control Handbook. Montreal Health Press, 2000.
2. Our Bodies, Ourselves. Boston Women's Health Book Collective, 2005.
3. Feminist Women's Health Center at www.FWHC.org and www.BirthControlComparison.info
4. Parenthood - Intrauterine Devices: national health care provider describes Paragard and Mirena as IUDs.
5. Petta C, Ferriani R, Abrao M, Hassan D, Rosa E Silva J, Podgaec S, Bahamondes L (2005). "Randomized clinical trial of a levonorgestrel-releasing intrauterine system and a depot GnRH analogue for the treatment of chronic pelvic pain in women with endometriosis." *Hum Reprod* 20 1993–8. doi:10.1093/humrep/deh869. PMID 15790607.
6. Marjoribanks J, Lethaby A, Farquhar C (2006). "Surgery versus medical therapy for heavy menstrual bleeding." *Cochrane Database Syst Rev*: CD003855. doi:10.1002/14651858.CD003855.pub2. PMID 16625593.
7. Faundes A, Alvarez F, Brache V, Tejada A (1988). "The role of the levonorgestrel intrauterine device in the prevention and treatment of iron deficiency anemia during fertility regulation." *Int J Gynaecol Obstet* 26 (3): 429–33. doi: 10.1016/00207292(88)90341-4. PMID 2900174.
8. IUDs—An Update. "Procedures for providing IUDs" <http://www.infoforhealth.org/pr/b6/B6boxes.shtml#procedures>.
9. "Understanding IUDs". Planned Parenthood. July 2005. <http://www.plannedparenthood.org/birth-control-pregnancy/birthcontrol/intrauterine-devices.htm>. Retrieved on 2006-10-08.
10. Bayer (2007). "Mirena U.S. Product Information". http://berlex.bayerhealthcare.com/html/products/pi/Mirena_PI.pdf. Retrieved on 2007-05-11. FDA Medical Review p.12
12. Ortiz M, Croxatto H (1987). "The mode of action of IUDs." *Contraception* 36 (1): 37–53. doi:10.1016/0010-7824(87)90060-6. PMID 3311625.
13. Stanford J, Mikolajczyk R (2002). "Mechanisms of action of intrauterine devices: update and estimation of postfertilization effects". *Am J Obstet Gynecol* 187 (6): 1699–708. doi:10.1067/mob.2002.128091. PMID 12501086., which cites:
14. Margret chandira, Debijit Bhowmik, Chiranjib, B jayakar , On intra-uterine drug delivery system - A review.
15. Chiranjib , Vinod Dubey, Debijit Bhowmik, k.k.tripathi, k.p.sampat kumar , Recent advances in intra-uterine drug delivery systems, *Annals of biological research* 2010 ,1,(1):70-75, www.scholarsresearchlibrary.com
16. Chinmaya keshari sahu, Amiyakanta Mishra, D. Venkata Ramana, Jimidi Bhaskar, Intra-uterine Devices- A review, volume-12, issue-3, 2020,
17. Zaida Urban Morlan, Luis Eduardo Serrona Mora, New developments in intra-uterine drug delivery system devices, <https://doi.org/10.1016/B978-0-12-819838-4.00018-3>.
18. Tapani Luukkainen , paivi pakarinen and juhani Toivonen, progestin- intra-uterine drug delivery systems, 12756C07 .PGS 10/22/01, 355-363.
19. Reed SD, Newton KM, Garcia RL, Allison KH, voigt JF, Jordan CD, et al 2010, complex hyperplasia with and without atypia: clinical outcomes and implications of progestin therapy, *American College of Obsterics and Gynaecology*, bol.106, no.4, pp.812-9.

20. Reed SD, Newton KM, Garcia RL, Allison KH, Voigt JF, Jordan CD, et al 2010, complex hyperplasia with and without atypia: clinical outcomes and implications of progestin therapy, American College of Obstetrics and Gynaecology, vol.106, no.4, pp.812-9.
21. Parveen parasar, Pinar Ozcan, Kathryan L. Terry, Endometriosis : Epidemiology Diagnosis and clinical management, Curr Obstet Gynecol Rep (2017) 6:34-41, DOI 10.1007/s13669-017-0187-1.
22. Stephan Gordts, Grigoris Grimbizis and Rudi campo, Symptoms and classification uterine adenomyosis, including the place of hysteroscopy in diagnosis, <https://www.researchgate.net/deref/https%3A%2F%2Fwww.fertsterdialog.com%2Fusers%2F16110-fertility-and-sterility%2Fposts%2F29040-25436>.