



Prevalence of Anaemia and Associated Risk Factors among Pregnant Women Attending Antenatal Care at General Hospital Dakin-Gari, Kebbi State, North West Nigeria.

Umar Garba Gwazawa¹, Sufiyanu Muhammad Dakin-gari², Mukhtar Shuaibu³, Abdullahi Ibrahim⁴

Department of Social Sciences, Kebbi State Polytechnic, Dakin-gari, Nigeria^{1,3}, Department of Mathematics, Kebbi State Polytechnic, Dakingari², Nigeria. Department of Public Health, National Open University Abuja⁴

ABSTRACT:

Anaemia is a global public health problem in both developing and developed countries; over 1.62 billion people suffer from anaemia, and is particularly more prevalent among pregnant women. The World Health Organization estimates that the prevalence of anaemia in pregnancy varies from 53.8% to 90.2% in developing countries and 8.3% to 23% in developed countries. This study is aimed to determine the prevalence of anaemia and its associated factors among pregnant women attending antenatal care clinic of General Hospital Dakin-Gari, Kebbi State, North west Nigeria. A total of 250 pregnant women were recruited from antenatal care clinic of General Hospital Dakin-Gari, Kebbi State using simple random technique after ethical approval was obtained from Research Ethics Committee of General Hospital Dakin-Gari, and consent from each study participant. Data on sociodemographics, economic status, and clinical history were collected using a pretested structured interviewer's questionnaire. Anaemia was diagnosed using the WHO-recommended cut-off of less than 33%. Participants were categorized into a subgroup of anaemia severity using the WHO cut-offs for mild, moderate, and severe anaemia. The overall prevalence of anaemia among pregnant women in this study was found to be 149 (59.6%). Of this, 113 (75.8%) had mild anaemia, while 36 (24.2%) had moderate anaemia. Residence, Marital status, monthly family income, occupation, Number of deliveries, Inter-pregnancy interval, and current malaria attack were the variables which had statistically significant association with the prevalence of anaemia. Based on the findings from our study, it showed a high prevalence of anaemia (59.6%) among pregnant women, making anaemia in pregnancy a severe public health problem. The predisposing risk factors associated with anaemia in this study were being in the rural area, having low monthly family income, women with less than eighteen months interval between pregnancies, women with greater than three number of deliveries and recent malaria attack. Educating pregnant women on anaemia should be intensified in antenatal care clinics.

KEYWORDS: Anaemia, Antenatal clinics, pregnant women, Prevalence

INTRODUCTION

Anaemia is a pathophysiological condition in which there is a marked reduction in the haemoglobin content of blood from the reference concentrations or in the number of red blood cells or defective maturation of red blood cells [1, 2]. It affects all age groups, but pregnant women and children are more vulnerable [2]. It is one of the most common nutritional deficiency diseases observed globally and affects more than 1.62 billion people of the world's population, of which 56 million are pregnant women. [3,4]. Anaemia is a major public health problem affecting all ages of the world's population, its highest prevalence being among pregnant women and children aged less than 5 years old. [4]. Anaemia in pregnancy affects about half of all pregnant women globally and constitutes an important global reproductive health issue [3]. The World Health Organization estimates that the prevalence of anaemia in pregnancy varies from 53.8% to 90.2% in developing countries and 8.3% to 23% in developed countries [4]. A haemoglobin concentration below 11.0g/dl or packed cell volume (PCV) of less than 33.0% is regarded as anaemia during pregnancy by the World Health Organization (WHO) [5]. Anaemia in pregnancy ranges from mild (10.0–10.9 g/L or haematocrit of 30–32.9%), moderate (7–9.9 g/L or 21–29.9%) to severe (less than 7 g/L or less than 21%) [6]. Iron deficiency, which results from a protracted negative iron balance, accounts for 50% of anaemia in women and 75% in women of reproductive age worldwide [7]. The lack of adequate dietary iron intake or absorption, the increased need for iron during pregnancy, and increased iron losses due to menstruation, worm infestation, and infections are all possible causes of the negative iron balance [8]. Anaemia in pregnant women continues to be one of the most intractable major public health issues, particularly in developing nations, because of a variety of sociocultural issues like a lack of essential nutrients like iron, folate, vitamin B12, C, and A, as well as a lack of awareness, poor dietary practices, parasitic infestations, blood loss, the Human Immunodeficiency Virus, tuberculosis, and malaria, as well as too early pregnancies, high parity, and poor uptake of ITNs among others [9]. Increased maternal age, reduced number of antenatal care visits, residing in rural areas, intestinal parasites, malaria parasitaemia, and being a primigravida are associated with anaemia in pregnancy [10]. Around the world, 500 million women of reproductive age suffer from anaemia. In 2011, 29% of non-pregnant women and 38% of pregnant women aged 15–49 years were anaemic worldwide, but the prevalence was highest in South Asia and Central and West Africa [11]. About 38.9% to 48.7% of pregnant women in South-East Asia, the Eastern Mediterranean, and Africa were anaemic [7]. In India, the

prevalence varies from 65% to 75% [12]. While studies across Nigeria reported anaemia prevalence rates ranging from 24.5% to 76.5% [13, 14]. The adverse effects of anaemia during pregnancy include general fatigue, foetal anaemia, low birth weight, preterm delivery, increased risk of postpartum haemorrhage, intrauterine growth restriction, perinatal mortality stillbirth, reduced work capacity, low infection tolerance, shortness of breath, and decreased physical and mental performance. The World Health Organization (WHO) advises daily iron and folic acid supplementation for pregnant women as part of antenatal care to prevent anaemia in pregnancy and intermittent iron and folic acid supplementation for menstruating women living in areas where the prevalence of anaemia is 20% or higher. It is believed that iron supplementation, routine deworming, consistent use of insecticide-treated bed nets, nutritional counselling, food fortification, iron and folic acid supplements, raising awareness of anaemia among pregnant women during their antenatal visits, raising awareness of the underlying causes and complications, and treating anaemia are vital to preventing anaemia and its complications [15,16]. It is believed that understanding the severity of anaemia and its contributing factors can help prevent harmful foetal and mother consequences. This study aimed to find out the prevalence and factors

MATERIALS AND METHODS

Study Region, Design, and Population.

The study was conducted at General Hospital Dakin-Gari, Kebbi State. General Hospital Dakin-Gari is a secondary health-care facility located in the Suru local government Headquarter Dakin -Gari , Kebbi State, North West, Nigeria. It serves as a referral centre in the Local government. The study was institution-based, descriptive, and cross-sectional in design and it included 250 pregnant women who attended the antenatal clinic of Sir Yahaya Memorial Hospital Birni-kebbi, from November 2022 to March 2023.

Sampling Technique.

The study participants were pregnant women attending antenatal care in General Hospital Dakin-Gari, Kebbi State. All pregnant women who consented to the study and who were not critically ill were eligible to participate in the study. The sample size was estimated at 250 using the formula for calculating the sample size for proportion in cross-sectional studies [17] a 75.7% prevalence of anaemia in pregnancy from a previous study [18] at 5% margin of error, and an anticipated 95% response rate. The eligible participants were selected by systematic sampling technique; one of five patients presenting consecutively at the clinic was enrolled into the study over a ninth clinic-day period until the required sample size was obtained.

Ethical Clearance.

The study design and methodology were reviewed by the ethical committee of General Hospital Dakin-Gari and approval was obtained. Written consent was obtained from consenting participants after the consent form was properly explained to them. Confidentiality was ensured as identifiers were not collected during data collection and access to the data collected was restricted to the research team only.

Data Collection and Laboratory Analysis.

A pretested structured interviewer's questionnaire was used to collect data on demographics and socioeconomic status by trained research assistants. About four millilitres (4ml) of venous blood was drawn into a syringe from each consenting participant and then transferred into a labelled EDTA container and transported to the laboratory for analysis. Trained laboratory technicians did the analyses in General Hospital Dakin-Gari, Kebbi State. PCV was estimated using a standard laboratory technique and procedure. Anaemia was diagnosed using the WHO-recommended cut-off packed cell volume (PCV) of less than 33%. Participants were categorized into a subgroup of anaemia severity using the WHO cut-offs for mild, moderate, and severe anaemia [6].

Data Management and Analysis.

Data was collected and entered into the database on a weekly basis after cleaning. Confidentiality of data collected was ensured as identifiers were not included and data access was strictly restricted to the research team only. At the end of the study, data was subject to descriptive and inferential statistical analysis using Statistical Package for Social Science (SPSS IBM Version 22.0) computer statistical software. Quantitative variables were summarized using mean and standard deviation, while qualitative variables were summarized using frequencies and percentages. Frequency distribution tables were constructed, and cross tabulations were done to examine the relationship between categorical variables. Chi-square analysis was used to determine the factors that were associated with anaemia. All levels of significance were set at $p < 0.05$.

RESULTS

Sociodemographic characteristics of respondents

All the 250 questionnaires administered were filled, returned and analysed, giving a response rate of 100%. The respondents' ages ranged from 18 to 42 years (mean = 26.56 ±4.89). A larger proportion of the respondents 127 (50.8%) were aged 21-29 years. most of the respondents 196 (78.4%) reside in rural areas, the majority of the respondents practised Islam as a religion 227 (90.8%), 223 (89.2%) of the 250 respondents were married, and most of them 196 (78.4%) belong to the Hausa ethnic group, and 120(48.0%) had Secondary education. A larger proportion of the respondents 155(62.0%) were full-time housewives, and 102(40.8%) respondents reported a monthly family income \geq N100, 000 as shown in [Table 1].

Obstetric and medical characteristics of respondents

Concerning the Obstetric and medical characteristics, majority of the respondents 112 (44.8%) were in the second trimester, 126 (50.4%) has 3 - 4 deliveries. 139 (55.6%) respondents has inter-pregnancy interval of >18 months and 201 (80.4%) has no history of Contraceptive use. 223 (89.2%) has no Vaginal bleeding during this pregnancy, 176(70.4%) respondents has history of Previous medical illness. 159 (63.6%) has malaria during Diagnosis of previous medical illness and 171 (68.4%) has no Current malarial attack as shown in [Table 2].

Table 1: Sociodemographic characteristics of respondents		
S/N	Variables	Frequency (%) n = 250
1.	Age (Years)	
	≤ 20	11 (4.4)
	21–29	127 (50.8)
	30–39	103 (41.2)
	> 40	9 (3.6)
2.	Residence	
	Rural	196(78.4)
	Urban	54(21.6)
3.	Religion	
	Muslim	227(90.8)
	Christian	23(9.2)
4.	Marital status	
	Married	223(89.2)
	Single	2(0.8)
	Widowed	6(2.4)
	Divorced/Separated	19(7.6)
5.	Tribe	
	Hausa	196(78.4)
	Fulani	23(9.2)
	Igbo	7(2.8)
	Yoruba	13(5.2)
	Others	11(4.4)
6.	Education level	
	Qur'anic education	12(4.8)
	Primary education	41(16.4)
	Secondary education	120(48.0)
	Tertiary education	77(30.8)
7.	Occupation	
	Housewife	155(62.0)
	Government employee	35(14.0)
	Private employee	26(10.4)
	Trader	13(5.2)
	Handicraft	21(8.4)
8.	Family monthly income (Naira)	
	≤50,000	57(22.8)
	51,000-99,000	91(36.4)
	≥100,000	102(40.8)

Table 2: Obstetric and medical characteristics of respondents		
SN.	Variables	Frequency (%) n = 250
1.	Duration of current pregnancy	
	First trimester	35(16.4)
	Second trimester	112(44.8)
	Third trimester	97(38.8)
2.	Number of deliveries	
	No births	11(4.4)
	1-2	84(33.6)

	3-4	126(50.4)
	>5	29(11.6)
3.	Inter-pregnancy interval	
	≤ 18 months	111(44.4)
	>18 months	139(55.6)
4.	Contraceptive use	
	Yes	49(19.6)
	No	201(80.4)
5.	Vaginal bleeding during this pregnancy	
	Yes	27(10.8)
	No	223(89.2)
6.	Previous medical illness	
	Yes	176(70.4)
	No	74(29.6)
7.	Diagnosis of previous medical illness	
	Malaria	159(63.6)
	Intestinal parasitosis	30(12.0)
	Others	61(24.4)
8.	Current malarial attack	
	Yes	79(31.6)
	No	171(68.4)

Table 3: Prevalence and pattern of anaemia among respondents

Variables	Frequency (%)
Haemoglobin concentration status (n = 250)	
Anaemic	149(59.6)
Non- Anaemic	101(40.4)
Pattern of anaemia (n = 149)	
Mild anaemia	113(75.8)
Moderate anaemia	36(24.2)

Table 4: Factors associated with anaemia among the respondents

Variables	Haemoglobin concentration status		Test of Significance
	Anaemic frequency (%)	Non Anaemic frequency (%)	
Age (Years)			
≤ 29	63 (45.7)	75 (54.3)	X ² = 2.5458 p-val = .4671
≥ 30	43 (51.8)	40 (48.2)	
Residence			
Rural	81 (41.3)	115(58.7)	X ² = 6.3917 p-val = .0115*
Urban	17 (68.0)	8(32.0)	
Religion			
Muslim	97(53.3)	85(46.7)	X ² = 3.8893 p-val = .0486*
Christian	14(35.9)	25(64.1)	
Marital status			
Married	121(60.2)	80(39.8)	X ² = 6.7869 p-val = .0092*
Others	6(30.0)	14(70.0)	
Tribe			
Hausa/Fulani	96(56.5)	74(43.5)	X ² = 5.5727 p-val = .2334
Other tribes	22(43.4)	29(56.6)	
Education level			
Primary and below	23(43.4)	30(56.6)	X ² = 1.6902 p-val = .6391
Formal	80(47.6)	88(52.4)	
Occupation			
Housewife	86(68.3)	40(31.7)	X ² = 18.21 p-val = .0011*
Employed	38(40.0)	57(60.0)	
Family monthly income (Naira)			
≤ 99,000	59(49.6)	60(50.4)	X ² = 7.1803

≥ 100,000	37(36.3)	65(63.7)	p-val = .0276*
Duration of current pregnancy			
First trimester	15(42.9)	20(57.1)	X ² = 5.071 p-val = .16667
Second trimester	47(56.6)	36(43.4)	
Third trimester	40(41.2)	57(58.8)	
Number of deliveries			
≤ 2	39(45.3)	47(54.7)	X ² = 0.3614 p-val = .0157*
≥ 3	79(58.5)	56(41.5)	
Contraceptive use			
Yes	13(44.8)	16(55.2)	X ² = 0.6101 p-val = .4348
No	101(52.6)	91(47.4)	
Vaginal bleeding during this pregnancy			
Yes	12(70.6)	5(29.4)	X ² = 2.0771 p-val = .1495
No	107(52.5)	97(47.5)	
Previous medical illness			
Yes	97(61.8)	60(38.2)	X ² = 0.1007 p-val = .7501
No	41(64.0)	23(36.0)	
Diagnosis of previous medical illness			
Malaria	93(66.9)	46(33.1)	X ² = 2.0513 p-val = .3586
Others	47(57.3)	35(42.7)	
Current malarial attack			
Yes	40(61.5)	25(38.5)	X ² = 15.7624 p-val = .000072*
No	51(32.7)	105(67.3)	

*Statistically significant (p < 0.05)

Prevalence, pattern and factors associated with anaemia among respondents

About 149 (59.6%) of the 250 pregnant women were anaemic. Of these, 113 (75.8%) had mild anaemia, while 36 (24.2%) had moderate anaemia and there is no severe anaemia found in this study as shown in [Table 3]. The proportion of respondents with anaemia was significantly higher (p < 0.05) among rural respondents (68.0%) as compared to those who resides in urban area (41.3%) and among those who are currently married (60.2%) as compared to those who are not currently married (30.0%). It was significantly higher among those who are fully house wives (68.3%) as compared to those who not fully house wives. It was also found significantly higher among respondents with monthly family income of ≤ 99,000 Naira (64.3%) as compared to those with ≥ 100,000 Naira (36.3%). It was also significantly higher among respondents who has ≥ 3 deliveries (58.5%) as compared to those with < 3 deliveries (45.3%) and among those with inter-pregnancy interval of ≤ 18 months (72.3%) as compared to those with > 18 months (25.7%). It was also found significantly higher among respondents with current malaria attack (61.5%) as compared to those with no malaria (32.7%). [Table 4].

DISCUSSION

The diagnosis of anaemia during booking among pregnant women is essential as it affords one the opportunity to institute interventions to prevent the complication of anaemia especially considering the prevalent high maternal and perinatal morbidity and mortality associated with anaemia in pregnancy in the tropics [19]. Data from the literature in developing countries have reported prevalence of anaemia in pregnancy that ranged from 35.0% to 75.0% [20]. This study analysed the prevalence and risk factors associated with anaemia among pregnant women attending the Antenatal Care Clinic of General Hospital Dakin-Gari, Kebbi State, Nigeria. The high prevalence of anaemia among the respondents in this study was (59.6%) which is in the middle range when compared to findings from other studies in Nigeria [17, 19, 20, 21,] and from countries in South Eastern Africa [22, 23].

In this study, the overall prevalence of anaemia among pregnant women was 59.6%. The high prevalence of anaemia in this study was mostly related to low sociodemographic factors and obstetrical history. The pattern of anaemia among the respondents in this study in which majority of them (75.8%) had mild anaemia, while the remaining ones (24.2%) had moderate anaemia, but none of them had severe anaemia which is in concordance with the findings in studies conducted in Enugu, South Eastern Nigeria in which most (90.7%) of the women had mild anaemia, a few of them (9.3%) had moderate anaemia, but none of them had severe anaemia [24]. Other studies across Nigeria generally reported similar results, with most respondents having mild anaemia, while a few had moderate anaemia, and very few or none of the respondents had severe anaemia [25, 26, 43]. In the present study, majority of anaemic patients belonged to the age group of 21–29 years (50.8%). It is comparable to the studies by [27,28,29] This may be due to the recurrent pregnancies and poor birth spacing in reproductive age women which make them more prone to anaemia in our study. Nearly 78.4% of anaemic subjects belonged to rural population while 21.6% in the urban population which is similar to (68%) and (65.9%) [30, 31]. The reason for higher burden of anaemia in the rural population may be related to inaccessibility of health-care centres. However, lack of awareness about the factors causing anaemia and possible strategies to prevent the risk factors of anaemia plays a vital role. Prevalence of anaemia was more in subjects (59.2%) having monthly income <10,000 per capital, this may be due to the reason that the low socioeconomic class was likely to be poorly educated and often has financial constraints. It is also supported by the observation that women with low income tend to consume diets that are low in micronutrients, animal protein, and vitamins [32]. It closely correlates with a studies by [33, 34, 35] where women of low socioeconomic group had higher prevalence of anaemia. When distributed according

to education levels, anaemia was found to be more prevalent in secondary level of education (48.0%). However, another study showed a direct relationship between impact of literacy and anaemia [36]. The present study was in concurrence with the other studies (37%) and (38.5%) who showed patients with anaemia in same group [37, 28]. In this study, the number of children were found to be significantly associated with anaemia in pregnant women ($p < 0.05$). The incidence of anaemia during pregnancy is expected to be higher as the number of pregnancies increases because of the repeated drain on the iron reserves. In fact, multiparity, especially when the pregnancies have occurred in a rapid sequence, is traditionally regarded as a cause of anaemia in pregnancy. Anaemia was more prevalent among the subjects who have family size >2 (62.0%) as compared to patients who have family size ≤ 2 (38.0%). There may be a direct relationship of family size with anaemia due to food insecurity for large family size; however, no significant relationship of anaemia with family size could be derived in the present study. Similarly, the higher incidence of anaemia in family size >2 was also reported [39].

In the present study, majority of anaemic respondents (44.4%) belonged to the second trimester as compared to the first and third trimesters which had (16.4%) and (38.8%) of patients, respectively. Similar results (36.5%) were reported in a study [26]. The physiological haemo-dilution that takes place during pregnancy reaches its peak during the second trimester and may account for the increase in anaemia seen at this time in pregnancy. Also the increased demand for micro-nutrient in pregnancy increases as the pregnancy progresses. It has been found in our study that (70.4%) of patients of anaemia were multigravida while 29.6% were primigravida. This may be explained by the fact that repeated childbirths make the patients more prone to malnutrition resulting in anaemia and reducing maternal iron reserves at every pregnancy by causing blood loss at each delivery. Other study also concluded that multiparous women had higher risk of anaemia [27]. Women with prior pregnancy sustain a 500–600 mg iron loss per pregnancy, which is increased by haemorrhage after delivery. Iron deficiency is, therefore, definitely more common as parity increases. About (55.6%) of respondents had anaemia with interpregnancy interval of > 18 months as compared to (44.4%) of respondents with interpregnancy interval of ≤ 18 months. On the contrary, higher prevalence of anaemia in those with repeated pregnancies with poor birth spacing was reported [41]. However, a cause-effect relationship still needs to be proved. In the present study, current malaria attack were found to be significantly associated with anaemia in pregnant women ($p < 0.05$), as were the findings of [42]. In order to avoid anaemia during pregnancy, receiving early antenatal care would serve as an important preventive measure.

CONCLUSION

Based on the findings from our study, it showed a high prevalence of anaemia (59.6%) among pregnant women attending the antenatal care clinic of General Hospital Dakin-Gari, Kebbi State, northwest Nigeria, making anaemia in pregnancy a severe public health problem. The predisposing risk factors associated with anaemia in this studies were being in the rural area, having low monthly family income, women with less than eighteen months interval between pregnancies, women with greater than three number of deliveries and recent malaria attack. Educating pregnant women on anaemia

ACKNOWLEDGMENTS

This research was supported by Tertiary Education Trust Fund (TETFund) under the 2022 Institutional Based Research (IBR) projects in Kebbi State Polytechnic Dakin-Gari, Kebbi State, Nigeria. We also wish to thank the management and staff of the General Hospital Dakin-Gari, Kebbi State, Nigeria.

REFERENCES

- [1]. Grewal A (2010) Anaemia and pregnancy: anaesthetic implications. *Indian Journal of Anaesthesia* 54,380–6.
- [2]. Lin L, Wei Y, Zhu W, Wang C, Su R, Feng H and Yang H (2018) Prevalence, risk factors and associated adverse pregnancy outcomes of Anaemia in Chinese pregnant women: a multicentre retrospective study. *BMC Pregnancy and Childbirth* 18, 111.
- [3]. R. Onoh, O. Lawani, P. Ezeonu, P. Nkwo, T. J. P. Onoh, and L. Ajah, "Predictors of anemia pregnancy among pregnant women accessing antenatal care in a poor resource setting in South Eastern Nigeria," *Sahel Medical Journal*, vol. 18, no. 4, pp. 182–187, 2015
- [4]. B. Bruno, E. Mclean, I. Egli, and M. Cogswell, *World Prevalence of Anaemia 1993–2005. WHO Global Database on Anaemia*, World Health Organisation, Geneva, Switzerland, 2008.
- [5]. R. I. Anorlu, A. A. Oluwole, and O. O. Abudu, "Sociodemographic factors in anaemia in pregnancy at booking in Lagos Nigeria," *Journal of Obstetrics and Gynaecology*, vol. 26, no. 8, pp. 773–776, 2006.
- [6]. S. Salhan, V. Tripathi, R. Singh, and H. S. Gaikwad, "Evaluation of haematological parameters in partial exchange and packed cell transfusion in treatment of severe anaemia in pregnancy," *Anaemia*, vol. 2012, Article ID 608658, 7 pages, 2012
- [7]. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *Lancet Glob Health*. 2013; 1(1):E16–25.
- [8]. Orish VN, Onyeabor OS, Boamong JN, Acquah S, Sanyaolu AO, Iriemenam NC. The effects of malaria and HIV co-infection on hemoglobin levels among pregnant women in Sekondi-Takoradi, Ghana. *Int J Gynecol Obstet*. 2013; 120(3): 236–9.
- [9]. Balarajan Y, Ramakrishnan U, Özaltın E, Shankar AH, Subramanian SV. Anaemia in low income and middle-income countries. *The Lancet* 2011; 378(9809): 2123–2135.

- [10]. Bassi AP, Idoko L, Dibigbo-Ibeaji NM, Adeniyi OG, Ramiyil MS, Ogundeko TO et al. Prevalence of anaemia in pregnancy among women visiting antenatal clinic in Bingham University Teaching Hospital Jos, Nigeria. *Clinical Medicine Research* 2016; 5(3): 52-62.
- [11]. Adam I, Ibrahim Y, Elhardello O. Prevalence, types and determinants of anemia among pregnant women in Sudan: a systematic review and metaanalysis. *BMC Hematol* 2018; 18: 31.
- [12]. Suryanarayana R, Chandrappa M, Santhuram AN, Prathima S, Sheela SR. Prospective study on prevalence of anemia of pregnant women and its outcome: a community-based study. *J Family Med Prim Care* 2017; 6(4): 739-43.
- [13]. Gwarzo MY, Ugwa EA. The pattern of anaemia in Northern Nigerian pregnant women. *J. Med. Med. Sci.* 2013; 4(8): 319-323.
- [14]. Idowu OA, Mafiana CF, Dapo S. Anaemia in pregnancy: a survey of pregnant women in Abeokuta, Nigeria. *Afr Health Sci* 2005; 5(4): 295-299.
- [15]. Morey SS. CDC issues guidelines for prevention, detection and treatment of iron deficiency. *Am Fam Physician* 1998; 58(6): 1475-7.
- [16]. World Health Organization (WHO). Strategies to prevent anaemia: recommendations from an expert group consultation, New Delhi, India, 5-6 December 2016. Available at: <https://apps.who.int/iris/handle/10665/312109> [Last accessed on 2022, September 17].
- [17]. Ibrahim T. Research Methodology and Dissertation Writing for Health and Allied Health Professionals. Abuja: Cress Global Link Limited; 2009.
- [18]. Isah BA, Ayotunde AS, Awosan KJ, Yunusa EU. Prevalence of anaemia in pregnancy and related factors among women attending antenatal care (ANC) in General Hospital, Bunza, Kebbi State, Nigeria. *Med Sci Healthcare Pract.* 2019; 3(2): 47-55.
- [19]. T. Agan, J. E. Ekabua, A. E. Udoh, E. I. Ekanem, E. E. Efiok, and M. A. Mgbekem, "Prevalence of anemia in women with asymptomatic malaria parasitemia at first antenatal care visit at the University of Calabar Teaching Hospital, Calabar, Nigeria," *International Journal of Women's Health*, vol. 2, no. 1, pp. 229–233, 2010.
- [20]. A. O. Omigbodun, "Recent trends in the management of anaemia in pregnancy," *Tropical Journal of Obstetrics and Gynaecology*, vol. 21, no. 1, pp. 1–3, 2004.
- [21]. R. I. Anorlu, A. A. Oluwole, and O. O. Abudu, "Sociodemographic factors in anaemia in pregnancy at booking in Lagos, Nigeria," *Journal of Obstetrics and Gynaecology*, vol. 26, no. 8, pp. 773–776, 2006.
- [22]. F. I. Buseri, E. K. Uko, Z. A. Jeremiah, and E. A. Usanga, "Prevalence and risk factors of anaemia among pregnant women in Nigeria," *Open Journal of Hematology*, vol. 2, pp. 14–19, 2008.
- [23]. J. O. Komolafe, O. Kuti, O. Oni, and B. E. Egbewale, "Sociodemographic characteristics of anaemic gravidae at booking: a preliminary study at Llesha, Western Nigeria," *Nigerian Journal of Medicine*, vol. 14, no. 2, pp. 151–154, 2005.
- [24]. N. R. van den Broek and E. A. Letsky, "Etiology of anemia in pregnancy in south Malawi," *The American Journal of Clinical Nutrition*, vol. 72, supplement 1, pp. S247–S256, 2000.
- [25]. A. K. M. Hogue, S. B. Kadar, E. Hogue, and C. Mugero, "Prevalence of anaemia in pregnancy at Greytown, South Africa," *Tropical Journal of Obstetrics and Gynaecology*, vol. 23, no. 1, pp. 3–7, 2006.
- [26]. Dim CC, Onah HE. The prevalence of anemia among pregnant women at booking in Enugu, South Eastern Nigeria. *Med Gen Med.* 2007; 9(3): 11.
- [27]. Idowu OA, Mafiana CF, Dapo S. Anaemia in pregnancy: a survey of pregnant women in Abeokuta, Nigeria. *Afr Health Sci* 2005; 5(4): 295-299.
- [28]. Isah BA, Ayotunde AS, Awosan KJ, Yunusa EU. Prevalence of anaemia in pregnancy and related factors among women attending antenatal care (ANC) in General Hospital, Bunza, Kebbi State, Nigeria. *Med Sci Healthcare Pract.* 2019; 3(2): 47-55.
- [29]. Ayano B, Amentie B. Assessment of prevalence and risk factors for anemia among pregnant mothers attending anc clinic at Adama hospital medical college, Adama, Ethiopia. *J Gynecol Obstet* 2018;6:31-9.
- [30]. Obai G, Odongo P, Wanyama R. Prevalence of anaemia and associated risk factors among pregnant women attending antenatal care in Gulu and Hoima regional hospitals in Uganda: A cross sectional study. *BMC Pregnancy Childbirth* 2016;16:76.
- [31]. Getahun W, Belachew T, Wolide AD. Burden and associated factors of anaemia among pregnant women attending antenatal care in southern Ethiopia: Cross sectional study. *BMC Res Notes* 2017;10:276.
- [32]. Mihiretie H, Fufa M, Mitiku A, Bacha C, Getahun D. Magnitude of anaemia and associated factors among pregnant women attending antenatal care in Nekemte health centre, Nekemte, Ethiopia. *J Med Microbiol Diagn* 2015;4:3.
- [33]. Weldemariam TB. Prevalence and factors associated with anemia among pregnant women attending antenatal care in Shallaworeda, Warsi zone, Oromia region. *Int J Green Pharm* 2018;12:253.
- [34]. Vander Jagt DJ, Brock HS, Melah GS, EI-Nafaty AU, Crossey MJ, Glew RH. Nutritional factors associated with anaemia in pregnant women in Northern Nigeria. *J Health Popul Nutr* 2007;25:75-81.

- [35]. Okube OT, Mirie W, Odhiambo E, Sabina W, Habtu M. Prevalence and factors associated with anaemia among pregnant women attending antenatal clinic in the second and third trimesters at Pumwani maternity hospital, Kenya. *J Obstet Gynecol* 2016;6:16-27.
- [36]. K, Fong Y, Arulkumaran S. Anaemia in pregnancy a cross-sectional study in Singapore. *Eur J Clin Nutr* 1998;52:65.
- [37]. Javed MT, Abbas N, Hussain R, Zahoor T. A study on iron deficiency anaemia and hematological differences around delivery in women of different socio-economic and age groups. *Med J Islam Acad Sci* 2001;14:151-60.
- [38]. Dutta PK, Nagraj T, Gopinath VP. A case control study of anemia in pregnancy. *Indian J Prev Soc Med* 1992;23:1-5.
- [39]. Yadav RK, Sawmy MK, Banjade B. Knowledge and practice of anaemia among pregnant women attending clinic in Dr. Prabhakar Kore hospital, Karnataka a cross sectional study. *J Dent Med Sci* 2014;13:74-80.
- [40]. Sharma P, Mehta S, Nagar R. Prevalence of anemia and socio-demographic factors associated with anemia among pregnant women attending antenatal hospital in Jaipur city, India. *Age (Omaha)* 2013;20:20-5.
- [41]. Gautam VP, Bansal Y, Taneja DK, Renuka S. Prevalence of anaemia amongst pregnant women and its socio-demographic associates in rural area of Delhi. *Indian J Community Med* 2002;27:157-60.
- [42]. Bello FZ, Awosan KJ, Umar AG, Danmadami AM, David EC, Lawal A. Prevalence and factors associated with anaemia among pregnant women attending the Antenatal Care Clinic of UDUTH, Sokoto, Nigeria. *Int Arch Med Health Res* 2022; 3(1): 1-9.
- [43]. Umar S, Sufiyanu MD, Abdullahi I. Prevalence of Anaemia and its associated factors among pregnant women attending antenatal care at Sir Yahaya Memorial Hospital Birnin Kebbi, North West Nigeria. *Int. Advance research journal in science, Engineering and Technology. Vol. 10, Issue 3, March 2023.*