



Mushroom Classification of Toxicity Using Deep Learning

*Mrs. Rajeshwari. P^{*1}, Harini K. R^{*2}, Harish Raghavan M.R^{*3}*

^{*1} Assistant Professor, Department of Software Systems and AIML, Sri Krishna Arts and Science College Coimbatore, Tamil Nadu, India

^{*2} PG Student of Computer Science, Sri Krishna Arts and Science College Coimbatore, Tamil Nadu, India

^{*3} PG Student of Computer Science, Sri Krishna Arts and Science College Coimbatore, Tamil Nadu, India

DOI : <https://doi.org/10.55248/gengpi.5.1024.2907>

ABSTRACT:

Mushroom classification has undergone significant advancements, evolving from traditional methods based on observable morphological traits to modern molecular techniques that offer greater accuracy and reliability. This paper explores the history and development of mushroom taxonomy, emphasizing the limitations of morphology-based approaches, particularly with cryptic species that are morphologically similar but genetically distinct. The shift towards molecular phylogenetics, especially DNA sequencing methods such as the Internal Transcribed Spacer (ITS) region, has revolutionized fungal classification, allowing for a more precise understanding of genetic relationships and the discovery of new species. This study also highlights the role of bioinformatics and genomic data integration in constructing phylogenetic trees, contributing to a comprehensive view of fungal biodiversity. The applications of accurate classification extend beyond taxonomy, impacting fields such as agriculture, medicine, biotechnology, and environmental conservation. The paper concludes by discussing the future of mushroom classification, with promising applications in bioremediation, industrial biotechnology, and medicinal mushroom development.

Keywords: Molecular phylogenetics, DNA sequencing, Cryptic species, Fungal biodiversity, Bioinformatics

1. INTRODUCTION

1.1 Overview of Mushroom Taxonomy

Mushroom taxonomy has evolved significantly over time, starting from traditional methods based on observable morphological traits such as size, shape, colour, spore formation, and reproductive structures. These features, though useful, had limitations due to the convergent evolution of some species. For instance, some mushroom species share very similar morphological characteristics but are not closely related. This is particularly problematic in large genera like *Agaricus* and *Amanita*, where there are many morphologically similar species.

Historically, mushrooms were classified based on physical traits and habitat, but this method often led to misidentification. For example, the *Agaricus* genus, which includes both edible and toxic species, was notoriously difficult to categorize using traditional methods alone. As a result, dangerous misidentifications occurred in foraging practices.

1.2 Modern Advances in Classification

With the advent of molecular biology, mushroom classification has shifted towards more accurate techniques involving DNA sequencing. Molecular methods such as ribosomal DNA (rDNA) sequencing (especially the Internal Transcribed Spacer or ITS region) have become the gold standard for species identification. These techniques enable researchers to look beyond physical similarities and understand the genetic relationships between fungi.

For instance, molecular techniques have helped distinguish between cryptic species—those that are morphologically indistinguishable but genetically distinct. In some cases, molecular analysis has led to the discovery of entirely new species. This shift to molecular classification has revolutionized fungal taxonomy and led to a more comprehensive understanding of mushroom biodiversity.

1.3 The Need for Accurate Classification

Accurate classification is crucial in fields such as agriculture, medicine, biotechnology, and ecology. Misidentifying mushroom species can have significant economic and health consequences. For instance, misidentifying a toxic mushroom as edible can lead to poisoning or death. In agriculture, understanding the specific types of mycorrhizal fungi that associate with crops can enhance crop yields and soil health.

Additionally, industrial applications of mushrooms—such as in biofuels and pharmaceutical compounds—depend on the accurate identification of species with desired traits. Proper classification helps industries to isolate and mass-produce these fungi for commercial purposes.

2. MOLECULAR PHYLOGENETIC AND GENOMICS

2.1 *The Basics of Molecular Phylogenetic*

Molecular phylogenetic uses genetic sequences to trace the evolutionary relationships between organisms. In fungi, the Internal Transcribed Spacer (ITS) region of ribosomal DNA has been widely adopted as a DNA barcode for identifying species. The ITS region is favoured because it evolves quickly and provides enough variation between species to make accurate distinctions.

For instance, ITS sequencing has been used to map out the phylogeny of the Basidiomycota, a large phylum that includes familiar mushrooms like *Agaricus Bosporus* (the common button mushroom). Using molecular data, researchers have constructed detailed phylogenetic trees that trace the evolutionary history of these organisms. This approach has revealed new insights into the relationships between different Basidiomycota species and their evolutionary pathways.

2.2 *Applications of Genomics in Mushroom Classification*

The development of whole-genome sequencing has enabled researchers to compare entire genomes between species, offering an even more precise understanding of genetic relationships. Whole-genome sequencing provides insights into the functional genes that drive species-specific traits such as toxin production, enzyme activity, and symbiotic relationships with plants. This technique also allows for the identification of genetic variations that contribute to adaptation and survival in different environments.

For example, genomic studies have revealed how certain mushrooms like *Pleurotus ostreatus* (oyster mushrooms) produce enzymes capable of breaking down lignin, a complex polymer in wood. This discovery has industrial applications, as these enzymes can be used in biofuel production and bioremediation of environmental pollutants.

2.3 *Case Study: Fungal Phylogenetic Tree of the Basidiomycota*

A landmark study in fungal phylogenetic used multi-locus sequencing (combining ITS, LSU, and rpb2 gene regions) to map the evolutionary history of the Basidiomycota. This study not only helped clarify relationships between well-known mushroom families but also identified several new cryptic species.

For instance, species in the genus *Coprinus* were split into several distinct genera, including *Coprinellus* and *Coprinopsis*, based on molecular evidence. This reclassification was significant for both scientific research and mushroom foraging, as it corrected years of misidentification.

3. CRYPTIC SPECIES AND INTEGRATIVE TAXONOMY

3.1 *Understanding Cryptic Species*

Cryptic species are morphologically similar or indistinguishable species that are genetically distinct. Traditional classification methods often failed to identify cryptic species, leading to an underestimation of biodiversity. Molecular techniques, such as multi-locus sequencing and DNA barcoding, have made it possible to distinguish these species.

For example, the *Lactarius* genus, which includes edible mushrooms prized by foragers, was thought to consist of a few widespread species. However, molecular analysis revealed

that many cryptic species exist within this genus, some with significantly different ecological roles and geographical distributions.

3.2 *Techniques in Integrative Taxonomy*

Integrative taxonomy combines data from multiple sources, including morphology, molecular genetics, ecology, and chemical analysis, to provide a more comprehensive classification system. This approach addresses the limitations of relying on a single data type, which can lead to misclassification.

An example of integrative taxonomy in practice is the reclassification of the *Boletus edulis* complex. These edible mushrooms, commonly known as porcini, were long thought to represent a single species. However, integrative studies combining ITS sequencing, ecological data, and chemical analysis revealed several distinct species within this complex, each with unique geographical and ecological niches.

3.3 Case Study: *Cryptic Species in the Boletus edulis Complex*

In a study conducted on the *Boletus edulis* complex, researchers found that what was once considered a single species consisted of multiple cryptic species with different ranges and ecological preferences. By using molecular data, researchers were able to clarify the taxonomy of this economically important group, leading to more accurate foraging practices and better understanding of their ecological roles. This revelation has also improved conservation strategies by identifying distinct species with specific habitat requirements. Additionally, it has enhanced the quality and safety of mushroom products in the market by ensuring more precise species identification.

4. ENVIRONMENTAL DNA (EDNA) AND METABARCODING

4.1 Introduction to Environmental DNA (eDNA)

Environmental DNA (eDNA) refers to genetic material that organisms leave behind in their environment, such as soil, water, or air. By extracting and analysing eDNA, researchers can detect the presence of species without physically capturing them. This technique is particularly useful in identifying fungi that are difficult to locate due to their cryptic nature or limited geographic range.

For example, eDNA has been used to detect rare mycorrhizal fungi in forest soils, where traditional surveying methods may fail to identify species that do not produce visible fruiting bodies. This non-invasive method expands our ability to monitor fungal biodiversity across large and inaccessible areas, providing crucial data for conservation efforts and ecological studies. Moreover, eDNA sampling helps track changes in fungal communities in response to environmental disturbances.

4.2 Metabarcoding Techniques for Mushroom Diversity

Metabarcoding is a high-throughput DNA sequencing method that allows researchers to identify multiple species from a single environmental sample. By sequencing the ITS region from environmental samples like soil or leaf litter, researchers can identify dozens or even hundreds of fungal species in a single analysis.

For instance, Metabarcoding has been used in forest ecosystems to catalogue fungal biodiversity, revealing species that were previously unknown or thought to be rare. This method has revolutionized fungal ecology by providing a more accurate picture of fungal communities and their roles in ecosystems.

4.3 Applications in Ecological Surveys

Ecological surveys using eDNA and Metabarcoding have uncovered new fungal species and provided valuable data for conservation efforts. For example, eDNA sampling in the Pacific Northwest revealed the presence of several rare and endangered fungi, some of which are critical for maintaining forest health. The ability to detect these species without disturbing the environment makes eDNA a powerful tool for ecological research.

5. BIOINFORMATICS AND DATA INTEGRATION IN MUSHROOM CLASSIFICATION

5.1 The Role of Bioinformatics in Fungal Taxonomy

The rise of bioinformatics has revolutionized the field of mushroom taxonomy by enabling researchers to analyse and compare vast amounts of genetic data. Databases like MycoBank, GenBank, and UNITE house millions of DNA sequences, providing a reference for identifying species based on their genetic material.

Bioinformatics tools like BLAST and PhyML allow researchers to compare genetic sequences across species and construct phylogenetic trees that represent evolutionary relationships. These tools are essential for analysing the large datasets generated by next-generation sequencing (NGS) techniques.

5.2 Integrating Molecular, Morphological, and Ecological Data

Data integration is a critical challenge in fungal taxonomy, as researchers must combine genetic, morphological, and ecological data to develop comprehensive classification systems. For example, molecular data may suggest that two species are closely related, but morphological or ecological differences could indicate that they occupy different niches or have distinct life histories.

The integration of multiple data types is particularly important in the classification of mycorrhizal fungi, which form symbiotic relationships with plants. By combining genetic data with information on the fungi's ecological roles, researchers can develop more accurate classifications that reflect both evolutionary history and ecological function.

5.3 Case Study: The UNITE Database

The UNITE database is a global repository for fungal ITS sequences, which are used to identify species.

By cataloguing sequences from around the world, UNITE helps researchers identify species that may be difficult to classify using traditional methods. For example, researchers studying fungal diversity in tropical rainforests have used UNITE to identify previously unknown species of ectomycorrhizal fungi, which play a crucial role in nutrient cycling. This database also facilitates large-scale ecological studies by allowing comparisons of fungal communities across diverse habitats, contributing to a deeper understanding of global fungal biodiversity.

6. PHYLOGENETIC NETWORKS AND HYBRID SPECIATION

6.1 Understanding Phylogenetic Networks

Traditional phylogenetic trees represent evolutionary relationships as a branching pattern, but they do not account for complex phenomena like horizontal gene transfer or hybridization. In fungi, where hybrid speciation is common, phylogenetic networks provide a more accurate representation of evolutionary history by showing how different lineages may have exchanged genes. These networks help uncover complex evolutionary pathways, offering insights into the genetic diversity and adaptation mechanisms of fungal species.

6.2 Hybrid Speciation in Fungi

Hybrid speciation occurs when two distinct species interbreed to form a new species with characteristics of both parents. In fungi, hybrid species are particularly common in ectomycorrhizal genera like *Russula* and *Lactarius*, where hybridization events have been documented using molecular data. These hybrid species can occupy unique ecological niches and contribute to fungal biodiversity, often exhibiting novel traits that enhance their adaptability to specific environmental conditions.

6.3 Case Study: Hybridization in *Pleurotus* (Oyster Mushrooms)

The *Pleurotus* genus, which includes popular edible mushrooms like *Pleurotus ostreatus* (oyster mushroom), is known for its ability to hybridize. Researchers have documented hybrid species within the genus using molecular markers, which help distinguish hybrid individuals from their parent species. This research has practical applications for mushroom cultivation, as hybrids often have desirable traits such as faster growth or higher yields.

7. FUTURE APPLICATIONS OF MUSHROOM CLASSIFICATION

7.1 Biotechnology and Industrial Applications

The accurate classification of mushrooms has significant implications for biotechnology. Many fungi produce enzymes, bioactive compounds, and secondary metabolites that have industrial applications. For instance, white-rot fungi like *Phanerochaete chrysosporium* produce lignin-degrading enzymes that are used in biofuel production and pulp and paper processing.

By accurately identifying species with these capabilities, industries can optimize the production of valuable enzymes and bioactive compounds. Molecular classification helps identify fungi with potential for bioremediation, where they are used to degrade environmental pollutants like heavy metals and hydrocarbons.

7.2 Medicinal Mushroom Development

Mushrooms are a rich source of medicinal compounds, including polysaccharides, terpenoids, and lectins, which have shown potential in treating diseases like cancer, diabetes, and cardiovascular diseases. Accurate classification is essential for identifying mushrooms with medicinal properties.

For example, the Reishi mushroom (*Ganoderma lucidum*) has long been used in traditional medicine, but molecular analysis has revealed that several distinct species within the genus *Ganoderma* are sold under the name "Reishi." Accurate classification ensures that consumers are getting the right product with the desired medicinal properties.

7.3 Agricultural and Environmental Sustainability

Mushrooms play a critical role in sustainable agriculture, particularly through their symbiotic relationships with plants. Mycorrhizal fungi, which form associations with plant roots, improve nutrient uptake and soil health. Accurate classification of mycorrhizal fungi is essential for optimizing their use in agriculture.

For example, arbuscular mycorrhizal fungi (AMF) form symbiotic relationships with crops like wheat, corn, and soybeans. By accurately classifying AMF species and understanding their ecological roles, farmers can use these fungi to enhance crop yields and reduce the need for chemical fertilizers.

7.4 Food and Nutritional Applications

As mushroom cultivation becomes more widespread, accurate classification is essential for identifying edible species and preventing the accidental harvest of toxic species. Many wild mushrooms are still foraged, and misidentification can lead to poisoning or death.

Accurate classification is also important for discovering new edible mushrooms with high nutritional value. For instance, wild mushrooms in Northeast China have been identified as rich sources of proteins, vitamins, and antioxidants. These species, previously unknown to science, have potential for commercialization in the global market.

7.5 Bioremediation and Environmental Clean-up

Mushrooms are increasingly being used for bioremediation, the process of using organisms to clean up environmental pollutants. Certain fungi can break down complex pollutants like pesticides, heavy metals, and hydrocarbons. By accurately classifying these fungi, researchers can identify the best species for different types of bioremediation projects.


For example, *Pleurotus ostreatus* has been shown to degrade polycyclic aromatic hydrocarbons (PAHs) in contaminated soils. Accurate classification of the species ensures that researchers are using the most effective strains for environmental clean-up.

7.6 Fungal Bioinformatics and Data Science

As bioinformatics and data science continue to evolve, they will play an increasingly important role in mushroom classification. Big data tools like machine learning are being used to analyze vast amounts of fungal genetic data, uncovering patterns that would be impossible to detect manually. Advances in computational methods are enhancing the accuracy of fungal taxonomy and phylogeny. This progress is paving the way for more precise identification of fungal biodiversity.

For example, machine learning algorithms applied to fungal genome sequences can predict species' metabolic capabilities, such as producing bioactive compounds or degrading pollutants. Databases like Fungi DB integrate genomic, transcriptomic, and proteomic data, enhancing analyses and insights into fungal functions and interactions.

8. SYSTEM ARCHITECTURE



9. CONCLUSION

The classification of mushrooms has significantly evolved from its origins in morphology-based taxonomy, with the advent of molecular techniques such as DNA barcoding and whole-genome sequencing revolutionizing the field. These advancements have enabled more accurate species identification and the discovery of cryptic species. Bioinformatics has further enhanced researchers' ability to manage large datasets, integrating molecular, morphological, and ecological data to provide a comprehensive understanding of fungal diversity. Looking ahead,

the future of mushroom classification holds great promise, with exciting applications in biotechnology, medicine, agriculture, and environmental sustainability. By refining classification systems and incorporating emerging technologies, researchers are poised to unlock the full potential of fungi for human use, including medicinal compounds, biofuels, and environmental cleanup efforts.

10. REFERENCES

- [1] Baldrian, P. (2016). *Fungal Diversity and Their Role in Soil Processes*. Springer, Berlin, Heidelberg.

-
- [2] White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. In *PCR Protocols: A Guide to Methods and Applications* (pp. 315-322). Academic Press, Inc.
- [3] Vilgalys, R. & Hester, M. (1990). Rapid Genetic Identification and Mapping of Enzymatically Amplified Ribosomal DNA from Several *Cryptococcus* Species. *Journal of Bacteriology*, 172(8), 4238-4246.
- [4] Hawksworth, D. L. (2001). The Magnitude of Fungal Diversity: The 1.5 Million Species Estimate Revisited. *Mycological Research*, 105(12), 1422-1432.
- [5] Hibbett, D. S., Binder, M., Bischoff, J. F., Blackwell, M., Cannon, P. F., Eriksson, O. E., ... Zhang, N. (2007). A Higher-Level Phylogenetic Classification of the Fungi. *Mycological Research*, 111(5), 509-547.
- [6] Ainsworth, G. C. (1973). *Introduction to the History of Mycology*. Cambridge University Press.
- [7] Binder, M., & Hibbett, D. S. (2006). Molecular Systematics and Biological Diversification of Fungi. *Mycological Research*, 110(6), 649-664.
- [8] Mueller, G. M., Schmit, J. P., Leacock, P. R., Buyck, B., Cifuentes, J., Desjardin, D. E., ... Wu, Q. (2007). Global Diversity and Distribution of Macrofungi. *Biodiversity and Conservation*, 16(1), 37-48.
- [9] Moncalvo, J. M., Lutzoni, F. M., Rehner, S.A., Johnson, J., & Vilgalys, R. (2000). Phylogenetic Relationships of Agaric Fungi Based on Nuclear Large Subunit Ribosomal DNA Sequences. *Systematic Biology*, 49(2), 278-305.
- [10] Peay, K. G., Kennedy, P. G., & Bruns, T. D. (2008). Fungal Community Ecology: A Hybrid Beast with a Molecular Master. *Bioscience*, 58(9), 799-810.
- [11] Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30(4), 772-780.
- [12] Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., & Chen, W. (2012). Nuclear Ribosomal Internal Transcribed Spacer (ITS) Region as a Universal DNA Barcode Marker for Fungi. *Proceedings of the National Academy of Sciences*, 109(16), 6241-6246.