



## On Finding Integer Solutions to Binary Cubic Equation $x^2 - xy = y^2 + y^3$

*J. Shanthi<sup>1\*</sup>, M. A. Gopalan<sup>2</sup>*

<sup>1</sup>Assistant Professor, Department of Mathematics, Shrimati Indira Gandhi College, Affiliated to Bharathidasan University, Trichy- 620 002, Tamil Nadu, India. Email: [shanthiharshi@gmail.com](mailto:shanthiharshi@gmail.com) , Orcid id :0009-0008-5945-410X

<sup>2</sup>Professor, Department of Mathematics, Shrimati Indira Gandhi College, Affiliated to Bharathidasan University, Trichy-620 002, Tamil Nadu, India. Email: [mayilgopalan@gmail.com](mailto:mayilgopalan@gmail.com), Orcid id :0000-0003-1307-2348

### ABSTRACT

The process of obtaining many integer solutions to non-homogeneous polynomial equation of degree three with two unknowns given by  $x^2 - xy = y^2 + y^3$  is illustrated. A few relations between the solutions are presented. A procedure for obtaining second order Ramanujan numbers through integer solutions of the given binary cubic equation is illustrated.

**KEYWORDS:** Binary cubic equation ,Non-homogeneous cubic equation ,Integer solutions ,Second order Ramanujan numbers

Notations

$$t_{m,n} = n \left[ 1 + \frac{(n-1)(m-2)}{2} \right]$$

$$P_n^5 = \frac{n^2(n+1)}{2}$$

### INTRODUCTION

The theory of Diophantine equations is an ancient subject that typically involves solving, polynomial equation in two or more variables or a system of polynomial equations with the number of unknowns greater than the number of equations, in integers and occupies a pivotal role in the region of mathematics. The subject of Diophantine equations has fascinated and inspired both amateurs and mathematicians alike and so they merit special recognition. Solving higher degree diophantine equations can be challenging as they involve finding integer solutions that satisfy the given polynomial equation. Learning about the various techniques to solve these higher power diophantine equation in successfully deriving their solutions help us understand how numbers work and their significance in different areas of mathematics and science. For the sake of clear understanding by the readers, one may refer the varieties of cubic Diophantine equations with multi variables [1-14]. It seems that much work has not been done regarding polynomial Diophantine equations of degree three with two unknowns. This paper aims at determining many integer solutions to non-homogeneous polynomial equation of degree three with two unknowns given by  $x^2 - xy = y^2 + y^3$ . A few relations between the solutions are presented. A procedure for obtaining second order Ramanujan numbers through integer solutions of the given binary cubic equation is illustrated.

### METHOD OF ANALYSIS

The non-homogeneous cubic equation with two unknowns under consideration is

$$x^2 - xy = y^2 + y^3 \quad (1)$$

Treating (1) as a quadratic in  $x$  and solving for the same, we have

$$x = \frac{y[1 \pm \sqrt{4y+5}]}{2} \quad (2)$$

Let

$$\alpha^2 = 4y + 5 \quad (3)$$

which, after some algebra, is satisfied by

$$y_0 = s(s+1) - 1, \alpha_0 = 2s + 1 \quad (4)$$

Assume the second solution to (3) as

$$\alpha_1 = h - \alpha_0, y_1 = h + y_0 \quad (5)$$

where  $h$  is an unknown to be determined. Substituting (5) in (3) and simplifying, we have

$$h = 2\alpha_0 + 4$$

and in view of (5), it is seen that

$$\alpha_1 = \alpha_0 + 4, y_1 = y_0 + 2\alpha_0 + 4$$

The repetition of the above process leads to the general solution to (3) as

$$\begin{aligned} \alpha_n &= \alpha_0 + 4n = 2s + 1 + 4n, \\ y_n &= y_0 + 2n\alpha_0 + 4n^2 = (s + 2n)(s + 2n + 1) - 1 \end{aligned} \quad (6)$$

From (2), we get

$$\begin{aligned} x_n &= \frac{y_n [1 \pm \alpha_n]}{2} \\ &= (2n + s + 1)y_n, -(2n + s)y_n \end{aligned}$$

Thus, we have two sets of integer solutions to (1) represented by

Set 1

$$\begin{aligned} x_n &= x_n(s) = (s + 2n + 1)[(s + 2n + 1)(s + 2n) - 1], \\ y_n &= y_n(s) = (s + 2n)(s + 2n + 1) - 1. \end{aligned}$$

Set 2

$$\begin{aligned} x_n &= x_n(s) = -(s + 2n)[(s + 2n + 1)(s + 2n) - 1], \\ y_n &= y_n(s) = (s + 2n)(s + 2n + 1) - 1. \end{aligned}$$

Considering Set 1, the following relations are observed:

$$(i) \quad y_{n+2}(s) - 2y_{n+1}(s) + y_n(s) = 8, n = 0, 1, 2, \dots$$

$$(ii) \quad y_n(s) + 1 = 2t_{3s+2n}$$

$$(iii) \quad x_n(s) - 2P_{s+2n}^5 + 1 \text{ is a perfect square}$$

$$(iv) \quad \frac{x_n(s)}{y_n(s)} \text{ is a perfect square when } s = k^2 n^2 + 2(k-1)n$$

- (v)  $\frac{x_n(s)}{y_n(s)} = 2 P_n^5$  when  $s = n^3 + (n-1)^2 - 2$
- (vi)  $y_n(s+2) - 2y_n(s+1) + y_n(s) = 2$
- (vii)  $y_n(s) [y_{n+1}(s) - y_n(s) - 2] = 4x_n(s)$
- (viii)  $y_n(s) [y_n(s+2) - y_n(s) - 2] = 4x_n(s)$
- (ix)  $y_n(s) [y_{n+2}(s) - y_n(s) - 12] = 8x_n(s)$
- (x)  $y_n(s)[y_n(s+1) - y_n(s)] = 2x_n(s)$
- (xi)  $\frac{x_n(s)}{y_n(s)} - s$  is a perfect square when  $n = 4 t_{3,N}$
- (xii)  $\left(\frac{x_n(s)}{y_n(s)} - s\right)^2 - 1 = 8 t_{3,n}$
- (xiii)  $\frac{x_n(s)}{y_n(s)} - 1 = t_{2k+4,n}$  when  $s = (k+1)n^2 - (k+2)n, k \geq 1$
- (xiv)  $\frac{x_n(s)}{y_n(s)} - 1 = t_{2k+3,n}$  when  $s = (2k+1)n^2 - (2k+1)n, k \geq 0$
- (xv)  $y_{n+1}(s) + y_n(s) = 2y_n(s+1) + 2$
- (xvi)  $y_{n+2}(s) + y_n(s) = 2y_n(s+2) + 8$
- (xvii)  $y_{n+1}(s) = y_n(s+2)$
- (xviii)  $y_{n+2}(s) + 3y_n(s) = 4y_n(s+1) + 12$
- (xix)  $y_{n+2}(s) - 2y_{n+1}(s) - 4y_n(s+2) + 8y_n(s+1) = 3y_n(s)$
- (xx) From the integer solutions to (1) given by Set 1, one may generate Second order Ramanujan numbers as shown below :

Illustration :

$$y_2(3) + 1 = 56 = 1 * 56 = 2 * 28 = 4 * 14 = 7 * 8$$

$$= F_1 \quad F_2 \quad F_3 \quad F_4$$

$$F_1 = F_2 \Rightarrow (56+1)^2 + (28-2)^2 = (56-1)^2 + (28+2)^2 \\ = 57^2 + 26^2 = 55^2 + 30^2 = 3925$$

$$F_1 = F_3 \Rightarrow (56+1)^2 + (14-4)^2 = (56-1)^2 + (14+4)^2 \\ = 57^2 + 10^2 = 55^2 + 18^2 = 3349$$

$$F_1 = F_4 \Rightarrow (56+1)^2 + (8-7)^2 = (56-1)^2 + (8+7)^2 \\ = 57^2 + 1^2 = 55^2 + 15^2 = 3250$$

$$F_2 = F_4 \Rightarrow (28+2)^2 + (8-7)^2 = (28-2)^2 + (8+7)^2 \\ = 30^2 + 1^2 = 26^2 + 15^2 = 901$$

$$F_3 = F_4 \Rightarrow (14+4)^2 + (8-7)^2 = (14-4)^2 + (8+7)^2 \\ = 18^2 + 1^2 = 10^2 + 15^2 = 325$$

Thus , 3925 ,3349 ,3250 ,901 ,325 represent second order Ramanujan numbers .

A similar observation may be performed by considering the solutions in Set 2.

## Conclusion

The polynomial equation of degree three with two unknowns has been studied to obtain non-zero integer solutions .The process of eliminating the square-root will be beneficial for the researchers. As cubic equations are plenty , one may attempt to determine the solutions in integers for other choices of cubic diophantine equations.

## References

- [1] M.A.Gopalan ,S.Vidhyalakshmi ,J.Shanthi ,On the Cubic Equation with Four Unknowns  $x^3 + 4z^3 = y^3 + 4w^3 + 6(x - y)^3$  ,International Journal of Mathematics Trends and Technology, Vol 20(1),Pg 75-84,2015
- [2]. M.A.Gopalan,S.Vidhyalakshmi,J.Shanthi , On ternary cubic Diophantine equation  $3(x^2+ y^2) - 5xy + x + y + 1 = 12z^3$ , International Journal of Applied Research,vol1,Issue-8,2015, 209-212
- [3] M.A.Gopalan ,S.Vidhyalakshmi ,J.Shanthi ,On Cubic Equation with Four Unknowns  $x^3 + y^3 + 2(x + y)(x + y + 2) = 19zw^2$ ,International Journal for Mathematics, Vol 2(3),Pg 1-8,2016
- [4] M.A.Gopalan ,S.Vidhyalakshmi ,J.Shanthi ,On The Non-homogeneous Cubic Equation with Five Unknowns  $9(x^3 - y^3) = z^3 - w^3 + 12p^2 + 16$ ,International Journal of Information Research and Review (IJIRR) , Vol 3(6),Pg 2525-2528,2016
- [5] M.A.Gopalan ,J.Shanthi ,On The Non-homogeneous Cubic Equation with Five Unknowns  $(a + 1)^2(x^3 - y^3) = (2a + 1)(z^3 - w^3) + 6a^2p^2 + 2a^2$ ,International Journal of Modern Sciences and Engineering Technology (IJMSET) Vol 3(5),Pg 32-36,2016
- [6] E.Premalatha, J.Shanthi, M.A.Gopalan On Non - Homogeneous Cubic Equation With Four Unknowns  $(x^2 + y^2) + 4(35z^2 - 4 - 35w^2) = 6xyz$ , Vol.14, Issue 5, March 2021, 126-129.
- [7] J.Shanthi,M.A.Gopalan, A search on Non -distinct Integer solutions to cubic Diophantine equation with four unknowns  $x^2 - xy + y^2 + 4w^2 = 8z^3$ , International Research Journal of Education and Technology,(IRJEdT), Volume2,Issue01, May 2021. 27- 32
- [8] S.Vidhyalakshmi,J.Shanthi,M.A.Gopalan,"On Homogeneous Cubic equation with fourUnknowns  $x^3 - y^3 = 4(w^3 - z^3) + 6(x - y)^3$  , International Journal of Engineering Technology Research and Management , 5(7) ,July 2021,180-185
- [9] S.Vidhyalakshmi,J.Shanthi,M.A.Gopalan, T. Mahalakshmi, " On the non-homogeneous Ternary Cubic Diophantine equation  $w^2 - z^2 + 2wx - 2zx = x^3$ , International Journal of Engineering Applied Science &Technology, July-2022, Vol-7, Issue-3, 120-121.
- [10] M.A. Gopalan, J. Shanthi, V.Anbuvali, Obervation on the paper entitled solutions of the homogeneous cubic eqution with six unknowns  $(w^2 + p^2 - z^2)(w - p) = (k^2 + 2)(x + y)R^2$ ,International Journal of Research Publication& Reviews, Feb-2023, Vol-4, Issue-2, 313-317.
- [11] J.Shanthi, S.Vidhyalakshmi,M.A.Gopalan, On Homogeneous Cubic Equation with Four Unknowns  $(x^3 + y^3) = 7zw^2$ ," Jananabha, May-2023, Vol-53(1), 165-172.
- [12] J. Shanthi , M.A. Gopalan, Cubic Diophantine equation is of the form  $Nxyz = w(xy + yz + zx)$  , International Journal of Modernization in Engineering Tech &Science, Sep-2023, Vol-5, Issue-9, 1462-1463.
- [13] J. Shanthi , M.A. Gopalan,"A Search on Integral Solutions to the Non- Homogeneous Ternary Cubic Equation  $ax^2 + by^2 = (a + b)z^3, a, b > 0$ ", International Journal of Advanced Research in Science, Communication and Technology, Vol-4, Issue-1, Nov-2024, 88-92.
- [14] J.Shanthi ,M.A.Gopalan ,On finding Integer Solutions to Binary Cubic Equation  $x^2 - xy = y^3$ ,International Journal of Multidisciplinary Research in Science ,Engineering and Technology ,,7(11),2024,16816-16820.