



Formulation and Evaluation of Tablet of Crude Extract of *Tinosporacordifolia* (Giloy) and *Cassia Angustifolia* (Senna)

¹AkashDivakar, Dr. Vijay Singh²

^{1,2} Department of Pharmacy, Six Sigma Institute of Technology and Science, Jafarpur Rudrapur
GILOY (*Tinosporacordifolia*)

ABSTRACT

Because natural products have no known adverse effects when compared to drugs, they are becoming more and more valuable in clinical research. According to traditional ayurvedic literature, *Tinosporacordifolia*, also known as "Guduchi," has extensive applications in the treatment of numerous ailments. Global interest in the plant has increased recently due to the identification of its active components and their biological role in disease prevention. The scope of our current research in this review includes

- (i) the genetic diversity of the plant and
- (ii) the biological function of the plant's isolated active ingredients in disease targeting.

Exploiting the biochemical and signaling pathways impacted by the compounds isolated from *Tinospora* remains the review's future focus.

KEY WORDS: Diabetes, Ayurveda, and natural products

1- INTRODUCTION:-

Tinosporacordifolia, commonly known as Amrita, Giloy, or Guduchi, is frequently used in Ayurvedic medicine's "Rasayanas" to boost the body's defenses against infections and strengthen the immune system [1]. This large, glabrous, deciduous climbing shrub, which belongs to the Menispermaceae family, is considered one of the most versatile and rejuvenating herbs and is used extensively in both folk and Ayurvedic medicine. The species is found in Thailand, Malaysia, Indonesia, and India. The plant's Hindi name, Giloy, is derived from a Hindu mythology that describes a heavenly elixir that celestial beings use to prevent aging and maintain their youthful appearance forever [2]. *T. cordifolia* has succulent stems that branch out into long, filiform, mushy aerial roots. The deeply left rosette-like bark is milky white to grey, resembling lenticels. The aerial portions and roots of *T. cordifolia* have yielded a vast number of chemicals that have been isolated. Yellow flowers emerge in clusters from nodes. When mature, drupes (fruits) turn crimson [3]. Numerous components, such as berberin, tinosporaside, tinosporin, tinocordifolioside, cordifolioside A, cordifolioside B, isocolumbin, and magnoflorine, have been isolated from various sections. Terpenoids, alkaloids, lignan, carbohydrates, bitters, steroids, and glycosides are all present. There have been discovered many components, including the glycoside giloin and the non-glucosides gilenin and gilosterol. The leaves include the alkaloids tinosporin, tinosporic acid, and tinosporol. A novel furanoid diterpene, tinosporide, was isolated from stems along with sitosterol, cordifol, heptacosanol, and octacosanol from leaves [4]. Berberin is a highly significant component found in the stem of *T. cordifolia*. It is an isoquinoline alkaloid with the chemical formula $C_{20}H_{18}NO_4$ and a molecular mass of 336.36122 g/mol. Under ultraviolet light, the yellow-colored alkaloid exhibits intense yellow fluorescence. It exhibits a variety of pharmacological activities that increase this plant's therapeutic efficacy [5].



FIGURE.01

History of *Tinosporacordifolia*:-

Giloy is a magical shrub that is used to treat skin and health issues. Did you know that in the past, Giloy was also known as Amrita? Amrita means "the nectar of life," and it is said that a full pot of amrita emerged from the ManthanduringSamudraManthan. The Devil and Demon started a battle for the amrita, and after the latter bit it and fled, Amrita dropped on the ground. These drops of Amrita are said to have developed into a climbing herb, called Giloy.

LatinName:

GiloyisalsoknownasTinosporacordifoliainLatin.

Family:

GiloyisamemberofthefamilyMenispermaceae.Itisaplantthatblossoms.Additionallyknownasmoonseedandcrescent,Menispermaceae.

Common Name:

ThemostpopularnamesforTinosporacordifoliaincludeGuduchi,Moonseed,andGiloya.

Habitat:

TropicalareasarefamiliartoGiloy.Itisatypeofvinethatgrowsbestindry,deciduousforestsandrequiresupport.

Season:

Thebesttimetogrowgiloyisfromspringtosummer.TheidealmonthsareJulyandAugustandFebruarytoMarch

2. ACTIVE CONSTITUENTS OF GILOY (6-7)

The giloy plant has a variety of active ingredients that are derived from its leaves, stems, roots, and other sections (Table 2).

3.THERAPEUTIC USES(8-15)

Tinosporacordifolia is used to cure a variety of ailments and has distinct ingredients. It is a multifunctional plant with multiple uses for its various dosage forms.

3.1 Immunity Enhancer

Giloy is used to strengthen or enhance immunity. Numerous antioxidants found in it combat free radicals, maintain the health of your cells, and eradicate illness. Giloy kills bacteria, purifies blood, and aids in the removal of poisons.

3.2 In Chronic Fever

Giloy aids in fever recovery. As an antipyretic medication, giloy can also lessen the symptoms of a number of potentially fatal illnesses, including malaria, swine flu, and dengue. In cases of fever, it enhances blood platelets.

3.3 In Digestion

Giloy is highly helpful in treating bowel-related problems and enhancing digestion. For best effects, take giloy powder with a little amla on a regular basis. It can also be used with jaggery to relieve constipation.

3.4 Treats Diabetes

Additionally, giloy is a hypoglycemic medication that aids in the treatment of diabetes, namely Type 2 diabetes. It aids in lowering blood sugar as well. It has been suggested that it mediates its ability to prevent diabetes by reducing oxidative stress (OS), encouraging insulin production, and blocking the processes of gluconeogenesis and glycogenolysis, which in turn controls blood sugar levels. *Tinosporacordifolia*'s main phytoconstituents include steroids, alkaloids, tannins, flavonoids, cardiac glycosides, and saponins; these compounds have been shown to have anti-diabetic properties.

3.5 Treats Arthritis

Anti-inflammatory and anti-arthritis qualities found in giloy aid in the treatment of arthritis and its various symptoms. thegiloy stem powder for arthritic conditions. It can be used to treat rheumatoid arthritis in conjunction with ginger. *Tinosporacordifolia* has the potential to be used as an anti-osteoporotic

agent because it has been shown to influence the mineralization, differentiation, and proliferation of bone-like matrix on osteoblast model systems in vitro.

3.6 Reduces Asthmatic Symptoms

Coughing, wheezing, chest tightness, and shortness of breath are all symptoms of asthma. Due to its anti-inflammatory properties, goly helps to lessen respiratory issues like tonsils, colds, and excessive coughing.

3.7 Improves Vision and reduces Signs of Aging

The giloy plant is used in several regions of India to improve vision clarity. To apply, bring giloy powder to a boil in water, allow it to cool, and then dab onto the eyelids. This herb has anti-aging qualities that help lessen wrinkles, fine lines, acne, and dark spots. It offers a person's skin to be smooth and radiant.

3.8 Anti-HIV effects

It has been demonstrated that TCE reduces the HIV virus's recurrent resistance, increasing the treatment result. TCE's anti-HIV effects were demonstrated by a decrease in the number of eosinophils, an increase in B cells, macrophages, polymorphonuclear leucocytes, and hemoglobin percentage, indicating its potential for use in the treatment of the illness.

SENNA(Cassia Angustifolia)

ABSTRACT

Sennaalata belongs to the Leguminosae family of medicinal herbs. It can be found in humid and tropical areas. Typhoid, diabetes, malaria, asthma, ringworms, tinea infections, scabies, blotch, herpes, and eczema are among the conditions for which the plant has historically been utilized. The purpose of the review is to provide light on *S. alata*'s pharmacological properties and ethnobotanical description. Folk medicine has documented the use of several plant components as medicinal agents to treat a range of illnesses and infections. The separated chemicals and extracts showed clear pharmacological activity. The array of secondary metabolites, including tannins, alkaloids, flavonoids, terpenes, anthraquinone, saponins, phenolics, cannabinoid alkaloids, 1,8-cineole, caryophyllene, limonene, α -selinene, β -caryophyllene, and others, may be responsible for the display of antibacterial, antioxidant, antifungal, dermatophytic, anticancer, hepatoprotective, antilipogenic, anticonvulsant, antihyperlipidemic, antimalarial, anthelmintic, and antiviral activities.

INTRODUCTION :-

growth of Mary's health care According to estimates from the World Health Organization (WHO), 80% of people living in underdeveloped nations primarily use herbal remedies. It was recently shown that about one-third of the medications that are regularly used come from natural sources. As a result, between 40,000 and 70,000 species of medicinal plants with exceptional therapeutic potential have been documented [16]. Approximately 200,000 phytochemicals were found when the pharmacological properties of herbal plants were evaluated scientifically. These substances support the purported therapeutic properties of plants and always provide evidence for the use of natural products in the creation of innovative medications. Notwithstanding these successes, the use of herbal remedies was not widely acknowledged in modern medicine since there was insufficient scientific proof and inadequate record-keeping. But the importance of herbs in pharmacology has made it necessary to provide scientific information on plant pharmacological tests and bioactive substances [17]. It is believed that medicinal plants, which have a wide range of biological uses, are the primary suppliers of chemicals having therapeutic potential. Finding these chemicals in medicinal plants typically leads to the isolation of new compounds and, ultimately, the creation of new drugs. Sennaalata is one of the medicinal plants having a variety of intriguing pharmacophores that have been the subject of scientific investigation. *S. alata*, sometimes called *Cassia alata*, is a herb belonging to the Leguminosae family that grows widely. It is also referred to as ringworm plant, candle bush, craw-craw plant, acapulo, or ringworm bush. The plant, which goes by various local names, is widely distributed throughout Asia and Africa. Numerous bioactive chemical substances are present in it. Phenolics (rhein, chrysaphanol, kaempferol, aloemodin, and glycosides), anthraquinones (alatinone and alatonal), fatty acids (oleic, palmitic, and linoleic acids), steroids, and terpenoids (sitosterol, stigmasterol, and campesterol) are a few of the chemical components that have been documented. Numerous biological functions are reportedly displayed by these secondary metabolites [18]. The flower, bark, seed, leaves, and root all showed a variety of biological activity. Antimicrobial, antifungal, anticryptococcus, antibacterial, anticancer, and anti-inflammatory properties are among these pharmacological actions [19]. antihelmintic, wound-healing, antioxidant, and antidiabetic properties. Drug-resistant illness outbreaks in recent years have resulted in a number of health problems. Pharmacological research has been directed toward the development of novel, effective, and safe medications derived from natural substances in an effort to address these problems. The ethnobotanical and pharmacological effects of *S. alata* are evaluated in this review, supporting the plant's various traditional uses[20].



FIGURE.02.



FIGURE.03.

HISTORY OF SENNA:-

With the probable exception of Aristotle, whose writings suggest a taxonomy, a real scientific attempt to categorize organisms did not take place until the 18th century, despite certain accounts of taxonomic history attempting to trace taxonomy to ancient civilizations. Previous publications were mostly descriptive and concentrated on plants with agricultural or medicinal value. This scientific way of thinking is divided into several levels. The so-called "artificial systems" that underpinned early taxonomy included Linnaeus's system of sexual classification for plants (the name "Systema Naturae" ("the System of Nature"), which suggested that Linnaeus thought his animal classification system was more than just a "artificial system"). Subsequently, "natural systems"—based on a more thorough analysis of taxonomic traits—arose. Examples of these systems are those of de Jussieu (1789), de Candolle (1813), and Bentham and Hooker (1862–1863). These categories were pre-evolutionary in their thinking and described empirical patterns.

After Charles Darwin's *On the Origin of Species* (1859) was published, a new theory of classifications based on evolutionary links was developed. From 1883 forward, this was the idea behind phyletic systems. The methods of Eichler (1883) and Engler (1886–1892) exemplified this technique.

When cladistic technique was developed in the 1970s, taxa were categorized using only the monophyly criterion—which was validated by the existence of synapomorphies. Since then, evidence from molecular genetics has been added to the body of knowledge, largely to the advantage of classical morphology [21].

CONCLUSION OF GILOY :-

A multipurpose plant is an asset to all living forms. Alkaloids, glycosides, lactones, and steroids are some of the active ingredients found in plant extracts. These active ingredients all play various physiological and immune-modulatory functions, highlighting the plant's wide range of adaptability. Research must be done on the ways that active substances genuinely interact with biological systems and influence the links between structure and function. We can identify new insights into our understanding of nature by examining the crystal structures of membrane-bound receptors, activating downstream signaling cascades, and seeing changes in the immediate environment at the site of action.

CONFLICT OF INTEREST

The authors affirm that there isn't any conflict of interest with this paper's publishing.

Table No. 1. Plant names in Indian languages

S. No.	Language	Name
	Sanskrit	Amrita
1.	Hindi	Giloy
2.	English	Tinospora
3.	Bengali	Golancha
4.	Gujarati	Gulvel
5.	Urdu	Gilo
6.	Punjabi	Gilow
7.	Telugu	Tippa-teega
8.	Odia	Guluchi
9.	Malayalam	Amrutu

Table No. 2. Active constituents of Giloy with Biological response

S. No.	Active Component Type	Compounds	Source	Biological Response
1.	Alkaloids	Berberine, Choline, Palmatine, Tembetarine, Magnofloicine, Tinosporin, Isocolumbin	Stem Root	Anticancer, Antiviral infections, Neurological Disorder and Anti-diabetic
2.	Glycosides	Tinocordiside, Cordioside	Stem	Treat Neurological Disorder like Parkinsons
3.	Terpenoid	Furanolactone	Whole plant	Vasorelaxants, Antiinflammatory, Antimicrobial, Antihypertensive, Antiviral
4.	Steroids	Beta-Sitosterol	Stem aerial parts	Induce Osteoporosis in early inflammatory arthritis
5.	Aliphatic compound	Octacosanol	Whole plant	Anti-nociceptive and anti-inflammatory
6.	Others	Glycin, Tinosporic acid	Root	Used to treat anxiety, Protease inhibitors for HIV

Conclusion of Senna:-

Senna plant has been essential in numerous other diseases; therefore, it may be employed to increase immunity in a pandemic infection. Senna has important nutritional components such as vitamins, minerals, phytochemicals, and antioxidants that can boost immunity and lower the risk of many diseases. Senna has several therapeutic properties, including antioxidants and laxatives that are useful in treating skin infections and constipation while also being beneficial against a variety of pathogens. All things considered, more thorough study and eventually clinical trials are required to investigate its medicinal potential on a commercial basis.

REFERENCE:-

- Warrier R.R, Singh B.G, Anandalakshmi R, Sivakumar V, Kumar A.M, Shivalingam R. "Vegetative Propagation of *Tinospora cordifolia* - A Folkloric and Ayurvedic Medicinal Plant", *Journal of Biomedical*, 2007; vol.2(2), pp: 131-137.
- Jalalpura S.S, Alagawadia K.R, Mahajanshetty C.S. "In Vitro Antihelmintic Property of Various Seed Oils," *Iranian Journal of Pharmaceutical Research*, 2006; vol.4, pp: 281-284.
- Singh B, Singh V.P, Kumar M. "Effect of *Tinospora cordifolia* Aqueous Extract on Traditional Food Crops of Garhwal Himalaya", *International Journal of Sustainable Agriculture*, 2009; vol.1(2), pp: 36-40
- Gupta R.S and Sharma A. "Antifertility Effect of *Tinospora cordifolia* (Willd.) Stem Extract in Male Rats", *Indian Journal of Experimental Biology*, 2004; vol.41(8), pp: 885-889.
- Mantena S.K, Sharma S.D, Katiyar S.K. "Berberin, A Natural Product Induces G-phase Cell Cycle Arrest and Caspase 3- Dependent Apoptosis in Human Prostate Carcinoma Cells", *Molecular Cancer Therapy*, 2006; vol.25(2), pp: 296-308.
- Sharmapriyavrat,Chaukambha Vishvbharti ,Ed-I,Re 2001 page.75
- Patel SS, Shah RS, Goyal RK. Antihyperglycemic, antihyperlipidemic and antioxidant effects of Dihar, a polyherbal ayurvedic formulation in streptozotocin induced diabetic rats. *Indian J Exp Biol*. 2009; 47: 564–70.
- Gupta R, Sharma V. Ameliorative effects of *Tinospora cordifolia* root extract on histopathological and biochemical changes induced by aflatoxin-b (1) in mice kidney. *Toxicol Int*. 2011; 18: 94–8
- Jagetia GC, Rao SK. Evaluation of the antineoplastic activity of guduchi (*Tinospora cordifolia*) in ehrlich ascites carcinoma bearing mice. *Biol Pharm Bull*. 2006; 29: 460–6
- Patel MB, Mishra S. Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. *Phytomedicine*. 2011; 18:1045–52

11. Ly PT, Singh S, Shaw CA. Novel environmental toxins: Steryl glycosides as a potential etiological factor for age-related neurodegenerative diseases. *J Neurosci Res.* 2007; 85:231–7.
12. Karpova EA, Voznyi Ya V, Dudukina TV, Tsvetkova IV. 4-Trifluoromethylumbelliferyl glycosides as new substrates for revealing diseases connected with hereditary deficiency of lysosome glycosidases. *Biochem Int.* 1991; 24:1135–44.
13. Kapil A, Sharma S. Immunopotentiating compounds from *Tinospora cordifolia*. *J Ethnopharmacol.* 1997; 58:89–95.
14. Chen S, Wu K, Knox R. Structure-function studies of DT-diaphorase (NQO1) and NRH: Quinone oxidoreductase (NQO2) *Free Radic Biol Med.* 2000; 29: 276–84.
15. Baldwin AS. Control of oncogenesis and cancer therapy resistance by the transcription factor NF-kappaB. *J Clin Invest.* 2001; 107: 241–6.
16. Oladeji O. S., Odelade K. A., Oloke J. K. Phytochemical screening and antimicrobial investigation of *Moringa oleifera* leaf extracts. *African Journal of Science, Technology, Innovation and Development.* 2019;12(1):79–84. doi: 10.1080/20421338.2019.1589082.
17. Svahn S. Uppsala, Sweden: Uppsala University; 2015. Analysis of secondary metabolites from *Aspergillus fumigatus* and *Penicillium nalgioense*; p. p. 195. Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Pharmacy.
18. Veeresham C. Natural products derived from plants as a source of drugs. *Journal of Advanced Pharmaceutical Technological Research.* 2012;3(4):200–201. doi: 10.4103/2231-4040.104709.
19. Makinde A. A., Igoli J. O., TA'Ama L., et al. Antimicrobial activity of *Cassia alata*. *African Journal of Biotechnology.* 2007;6(13):1509–1510.
20. Varghese G. K., Bose L. V., Habtemariam S. Antidiabetic components of *Cassia alata* leaves: identification through α -glucosidase inhibition studies. *Pharmaceutical Biology.* 2013;51(3):345–349. doi: 10.3109/13880209.2012.729066.
21. [https://en.m.wikipedia.org/wiki/Taxonomy_\(biology\)](https://en.m.wikipedia.org/wiki/Taxonomy_(biology))