WWW.IJRPR.COM

International Journal of Research Publication and Reviews

Journal homepage: www.ijrpr.com ISSN 2582-7421

Efficacy of Virtual Reality Immersive Gaming Combined with Dual-Task Gait Training in Chronic Post-Stroke Patients with Cognitive-Motor Interference

Dr. Diwakar Upadhyay, BPT ,MPT

MahaYogi Gorakhnath University Gorakhpur

ABSTRACT :

Background: Cognitive-motor interference (CMI) presents a major clinical challenge in chronic stroke recovery, often limiting community ambulation and increasing fall risks. Traditional rehabilitation protocols tend to target motor execution in isolation, largely overlooking the cognitive demands inherent to real-world gait. **Objective:** This framework evaluates the therapeutic potential of combining fully immersive Virtual Reality (VR) gaming with dual-task gait protocols to improve gait performance, dynamic stability, and executive integration in chronic post-stroke individuals experiencing CMI. **Methodology:** We outline a prospective, assessor-blinded randomized controlled trial design comparing a specialized VR-based dual-task protocol against conventional dual-task gait training over an eight-week intervention period. Primary evaluations focus on dual-task cost (DTC), spatiotemporal gait parameters, and task-switching capacity. **Conclusion:** Blending immersive 3D environments with targeted cognitive-motor gait tasks leverages neuroplastic principles to enhance attentional resource allocation, ultimately facilitating safer and more autonomous functional mobility.

Index Terms: *Stroke Rehabilitation; Virtual Reality; Dual-Task Training; Gait Analysis; Cognitive-Motor Interference; Executive Function; Neuroplasticity.*

1. Introduction

Despite advancements in acute stroke management, long-term functional recovery remains incomplete for many survivors. Persistent gait deficits frequently compromise independence, with **Cognitive-Motor Interference (CMI)** standing out as a particularly debilitating obstacle. CMI occurs when the simultaneous execution of a motor task and a cognitive operation leads to performance decrements in one or both domains.

In daily life, locomotion is rarely an isolated action. Navigating crowded sidewalks, crossing street intersections, or conversing while walking requires continuous executive oversight and ongoing cognitive adaptability. In chronic stroke survivors, damaged neural pathways and diminished executive reserve force individuals to dedicate heightened conscious effort to basic mechanics. When exposed to secondary cognitive demands, these patients exhibit slowed walking speeds, increased step-time variability, and heightened vulnerability to loss of balance.

While traditional physical therapy emphasizes isolated motor repetitions or basic dual-task drills (such as counting while walking), these approaches often lack the multisensory engagement and real-world complexity needed to drive robust cortical reorganization. Immersive Virtual Reality (VR) gaming bridges this gap. By embedding patients within dynamic, interactive 3D spaces, VR requires real-time perceptual processing and cognitive decision-making during active locomotion. This paper explores the combined efficacy of immersive VR gaming and dual-task gait protocols as an advanced therapeutic model for CMI in chronic stroke rehabilitation.

2. Neurobiological Mechanisms

The rationale for integrating immersive VR gaming into dual-task locomotion relies on interconnected neuroplastic mechanisms:

- **Enriched Environments and Cortical Remodeling:** Immersive VR delivers synchronized visual, auditory, and kinesthetic inputs. This continuous feedback stimulates premotor and sensorimotor networks, encouraging functional reorganization within perilesional pathways and supporting cortical remapping.
- **Automation of Gait Control:** By challenging central attentional resources in a controlled setting, dual-task VR protocols help transition movement control from conscious, energy-intensive compensations toward more automatic gait execution.
- **Dopaminergic Modulation via Gamification:** The goal-driven nature of interactive gaming—incorporating immediate performance feedback and achievement markers—promotes central dopamine release. Elevated dopaminergic activity enhances long-term potentiation (LTP) within motor circuits, facilitating skill retention and learning consolidation.

3. Proposed Methodology

Study Design & Participant Criteria

A randomized, assessor-blinded controlled trial is designed to evaluate clinical outcomes. Eligible participants must meet the following criteria:

1. Diagnosis of chronic stroke (>6 months post-onset).
2. Capacity to walk independently for at least 10 meters, with or without assistive devices.
3. Documented CMI, demonstrated by a measurable performance decrement during baseline dual-task testing.
4. Absence of uncorrected visual impairments, severe vestibular dysfunction, or active seizure disorders.

Rehabilitation Protocols

- **Intervention Group (VR Dual-Task):** Participants complete 45-minute sessions, three times weekly for eight weeks. Utilizing a head-mounted display (HMD) or immersive projection system, patients perform interactive tasks—such as avoiding virtual obstacles, selecting targets, or solving spatial puzzles—while walking on a instrumented treadmill or overground pathway.
- **Control Group (Conventional Dual-Task):** Participants follow an identical schedule receiving traditional dual-task training, consisting of overground walking paired with standard cognitive tasks (e.g., verbal fluency or serial subtraction) without virtual environments.

4. Primary Outcome Measures

Domain	Assessment Tool	Clinical Target
Gait Performance	10-Meter Walk Test (10MWT) & Motion Sensors	Measures comfortable walking speed and spatiotemporal symmetry under single and dual-task conditions.
Cognitive-Motor Cost	Dual-Task Cost (\% \text{DTC})	Quantifies relative degradation in walking speed when a cognitive task is introduced.
Dynamic Balance	Functional Gait Assessment (FGA) / TUG-Dual Task	Assesses balance control, direction changes, and stability during complex mobility tasks.
Executive Function	Trail Making Test (Part B) / Stroop Color-Word Test	Evaluates cognitive flexibility, working memory, and response inhibition speed.

\text{Dual-Task Cost (DTC)} = \left(\frac{\text{Single-Task Gait Speed} - \text{Dual-Task Gait Speed}}{\text{Single-Task Gait Speed}} \right) \times 100

5. Discussion & Clinical Implications

Combining immersive VR environments with dual-task paradigms represents an important evolution in neurological rehabilitation. Addressing CMI through interactive technology targets both physical and cognitive components of functional mobility simultaneously.

A primary advantage of this approach is ecological validity. Virtual scenarios can safely simulate real-world challenges—such as busy urban crosswalks or obstacles—allowing clinicians to train cognitive-motor adaptability without exposing patients to physical risk. Additionally, motion sensors integrated with VR platforms capture objective, precise spatial-temporal metrics, enabling tailored progressions based on individual performance. Finally, the interactive nature of gamified rehabilitation enhances patient motivation and session adherence, which are crucial factors for driving long-term neuroplastic adaptation.

6. Conclusion

Cognitive-motor interference remains a key contributor to mobility limitations and fall risks in chronic stroke survivors. Integrating immersive VR gaming with structured dual-task gait protocols offers an evidence-based method to promote neuroplasticity, restore gait automaticity, and improve functional independence. Future large-scale randomized clinical trials will be valuable for establishing standardized treatment parameters and evaluating long-term skill retention.