WWW.IJRPR.COM

International Journal of Research Publication and Reviews

Journal Homepage: www.ijrpr.com ISSN 2582-7421

A Review Study on Herbal Azadirachta Indica

Shivpal[1], Dr. Revathi A. Gupta[2], Mrs. Priyanka Yadu [3]

[1]Research Scholar, Pharmacy, Institute of Pharmacy, Dr. APJ Abdul Kalam University, Indore, M.P, India.
[2]Principal, Pharmacy, Institute of Pharmacy, Dr. APJ Abdul Kalam University, Indore, M.P, India.
[3]Assistant Professor, Pharmacy, Institute of Pharmacy, Dr. APJ Abdul Kalam University, Indore, M.P, India.

ABSTRACT:

Azadirachta indica commonly known as neem, is a medicinal tree belonging to the family Meliaceae. It has been widely used in traditional Indian medicine because of its diverse biological properties and the availability of bioactive constituents in different parts of the plant. Leaves, bark, seeds, flowers, roots and fruits have been investigated for their potential medicinal and industrial applications. Neem contains several classes of phytochemicals, including limonoids, flavonoids, tannins, phenolic compounds and other secondary metabolites. Among its important constituents are azadirachtin, nimbin, nimbolide and gedunin. Scientific investigations have reported antioxidant, antimicrobial, anti-inflammatory, antidiabetic and other pharmacological activities associated with neem extracts and isolated compounds. In addition to its medicinal importance, neem is widely recognized for its insecticidal properties and applications in agricultural pest management. However, the biological effects of neem depend on factors such as the plant part used, extraction method, concentration and chemical composition of the preparation. Although experimental findings indicate considerable therapeutic potential, further research is required to establish appropriate doses, long-term safety and clinical effectiveness in humans. This review discusses the botanical characteristics, phytochemical composition, traditional applications, pharmacological activities, safety considerations and future prospects of Azadirachta indica as a potential source of natural therapeutic agents

Keywords: Azadirachta indica, Neem, Phytochemicals, Herbal medicine, Pharmacological activities, Antioxidant, Antimicrobial, Medicinal plants.

1. Introduction:

Medicinal plants have played an important role in healthcare systems for centuries. Their use is based on traditional knowledge, observations of biological effects and the discovery of naturally occurring substances with potential therapeutic applications. Even with the development of modern pharmaceutical technology, plants continue to attract scientific interest as sources of new chemical compounds, research leads and ingredients for healthcare products. Investigations of medicinal plants also provide opportunities to understand the chemical basis of traditional remedies and to identify compounds that may be useful in future drug development. Azadirachta indica A. Juss., commonly called neem, is an important medicinal tree belonging to the family Meliaceae. It is widely cultivated in India and other tropical and subtropical regions. Neem is particularly valued for its adaptability to warm climates, multiple practical uses and extensive history in traditional medicine. Different parts of the tree have been used in traditional preparations for skin problems, oral hygiene, fever and other health complaints. Its leaves, bark, seeds, flowers and seed oil have also attracted attention in pharmaceutical, cosmetic and agricultural research [1,2].

The medicinal importance of neem is closely associated with its complex phytochemical composition. Investigations have identified several groups of secondary metabolites, including limonoids, flavonoids, tannins and phenolic compounds. Azadirachtin is particularly well known for its insecticidal activity, whereas compounds such as nimbolide, nimbin and gedunin have been investigated for various biological effects. These constituents do not necessarily occur in the same concentrations in every part of the plant. Their abundance and activity may vary with geographical origin, environmental conditions, maturity and extraction procedures [1–3].

Several experimental studies have examined the pharmacological properties of neem extracts and individual compounds. Reported activities include antimicrobial, antioxidant, anti-inflammatory and antidiabetic effects. Research has also investigated neem-derived substances in relation to cellular signalling, oxidative stress and cancer-related biological processes. These findings have increased scientific interest in neem as a possible source of pharmaceutical research leads [2,3]. Nevertheless, results obtained using isolated compounds, laboratory assays or animal models cannot automatically be interpreted as proof of therapeutic effectiveness in humans. The study of neem is also relevant to pharmaceutical formulation and quality control. Herbal preparations may contain multiple constituents that interact with one another, and their chemical composition can differ according to the method of preparation. Therefore, correct botanical identification, suitable extraction procedures, phytochemical characterization and appropriate safety assessment are important for developing reliable neem-based products. The present review summarizes the botanical characteristics, traditional uses, major phytochemical constituents and reported pharmacological activities of Azadirachta indica. It also considers its safety limitations and the research requirements that must be addressed before its medicinal potential can be translated into evidence-based healthcare applications

2. Botanical Profile Of Azadirachta Indica:

2.1 Taxonomical Classification

The taxonomical classification of neem is presented in Table 1.

Table 1. Taxonomical classification of *Azadirachta indica*

Taxonomic category	**Classification**
Kingdom	Plantae
Division	Tracheophyta
Class	Magnoliopsida
Order	Sapindales
Family	Meliaceae
Genus	*Azadirachta*
Species	*Azadirachta indica* A. Juss.
Common name	Neem

2.2 Morphological Characteristics

Neem is an evergreen to semi-evergreen tree, although it may shed some of its leaves under unfavourable environmental conditions. It generally develops a broad crown with numerous branches and can grow to a considerable height under suitable conditions. The bark is commonly greyish-brown and becomes rougher as the tree matures.

The leaves are alternate and pinnately compound, with several leaflets arranged along a central axis. Individual leaflets are generally elongated, with serrated margins and a pointed tip. Small, fragrant, whitish flowers occur in branched clusters. The fruit is a smooth, oval or ellipsoidal drupe that changes from green to yellowish when ripe. It usually contains a seed surrounded by a fleshy outer layer.

Different plant parts have distinct chemical compositions and uses. Leaves and bark are commonly investigated for their phenolic constituents and biological activities, while seeds are important sources of neem oil and limonoids. These differences should be considered when selecting plant material for research or formulation development [1,2].

2.3 Geographical Distribution and Habitat

Neem is widely associated with the Indian subcontinent and is cultivated across many tropical and subtropical regions. It grows well in warm environments and is known for its ability to tolerate relatively dry conditions. The tree is commonly found in agricultural areas, roadsides, gardens and other open locations.

Its adaptability makes neem useful for shade, soil protection and various agricultural purposes. However, growth conditions and environmental stress can influence the chemical composition of the plant. Consequently, plant material collected from different regions may not provide identical phytochemical profiles or biological responses.

3. Traditional and Medicinal Uses:

Neem has a long history of use in traditional Indian healthcare practices. Different parts of the tree have been incorporated into preparations intended for oral hygiene, skin care and the management of various common health complaints. The leaves, bark, seed oil and other plant materials have also been used in household preparations and traditional formulations [1,2].

Neem leaves are commonly associated with traditional skin-care practices. Preparations containing neem-derived ingredients have been used externally, although the benefits and safety of a particular preparation depend on its composition, concentration and method of application.

Neem twigs have traditionally been used for cleaning teeth. This practice has encouraged scientific interest in neem-containing oral-care products and their possible effects on microorganisms associated with oral hygiene. Neem extracts and related preparations have also been investigated for potential applications in toothpaste, mouthwash and other personal-care products.

Neem seed oil is another important plant-derived material. In addition to traditional external applications, it is used in agricultural formulations because of its effects on insects and other pests. Its composition differs from that of aqueous or alcoholic leaf extracts, and its safety cannot be assumed to be identical to that of other neem preparations.

Although traditional use provides useful information for selecting plants for scientific investigation, it does not independently establish clinical efficacy. Each proposed therapeutic application requires suitable evidence concerning effectiveness, dosage, formulation and safety.

4. Phytochemical Constituents Of Azadirachta Indica:

Phytochemical constituents are naturally occurring compounds that contribute to the chemical and biological properties of medicinal plants. Neem contains a diverse range of secondary metabolites, and the distribution of these substances differs among its leaves, bark, flowers, fruits and seeds. Several constituents have been isolated and studied to understand their potential biological activities [1–3].

Table 2. Major phytochemical groups reported in neem and their research significance

Phytochemical group	Representative constituents	Research significance
Limonoids	Azadirachtin, nimbolide, nimbin, gedunin	Insecticidal and other investigated biological activities
Flavonoids	Quercetin and related flavonoids reported in neem	Potential antioxidant and anti-inflammatory effects
Phenolic compounds	Various phenolic constituents	Antioxidant activity in experimental assays
Tannins	Different tannin-type compounds	Astringent and investigated antimicrobial effects
Sterols	Plant sterols	Chemical and biological significance
Fatty acids	Fatty acids present in seed oil	Properties relevant to oils and formulation development

Note: The presence and concentration of individual constituents depend on the plant part, geographical source and extraction method.

5. Pharmacological Activities:

5.1 Antimicrobial Activity

Antimicrobial activity is one of the widely investigated properties of neem. Extracts prepared from different plant parts have been studied against selected bacterial and fungal organisms. The results of these investigations vary according to the microorganisms tested, the type of extract, the concentration used and the experimental method [1,2].

The observed effects may be associated with multiple phytochemicals rather than a single active substance. Depending on the extract, these compounds may affect microbial growth, cellular membranes or other biological processes. However, the presence of an inhibition zone in an agar diffusion assay does not establish that an extract will be effective as a medicine in humans.

Further studies should determine minimum inhibitory concentrations, minimum bactericidal or fungicidal concentrations where appropriate, reproducibility and activity against clinically relevant isolates. Comparisons with established antimicrobial agents and investigations of toxicity are also necessary before therapeutic claims can be justified.

5.2 Antioxidant Activity

Oxidative stress occurs when the production of reactive species exceeds the capacity of biological antioxidant systems to maintain cellular balance. Excessive oxidative stress can damage cellular components, including lipids, proteins and nucleic acids.

Neem extracts have been investigated for antioxidant activity in several laboratory models. Phenolic compounds and flavonoids may contribute to the scavenging of free radicals under particular experimental conditions. Research has also examined the effects of neem-derived compounds on oxidative stress-related pathways [2,3].

Common antioxidant assays measure different chemical reactions, so results from one method may not directly correspond to results from another. In addition, antioxidant activity in a test tube does not necessarily indicate that a preparation will produce the same effect in the human body. Absorption, metabolism, tissue distribution and achievable concentrations must be considered.

5.3 Anti-inflammatory Activity

Inflammation is a biological response that helps the body react to injury and infection. Although it is essential for normal defence and tissue repair, excessive or persistent inflammation can contribute to disease.

Neem extracts and selected constituents have been studied for their possible effects on inflammatory mediators and cellular signalling pathways. Experimental investigations have reported changes in certain inflammatory responses, including pathways associated with the production of inflammatory molecules [2,3].

These findings provide a basis for further pharmacological research. Nevertheless, the magnitude and relevance of the effects depend on the specific compound, experimental model and dose. More controlled studies are required to establish the effectiveness and safety of standardized neem preparations for inflammatory conditions.

5.4 Antidiabetic Activity

Diabetes mellitus is a metabolic disorder characterized by persistent abnormalities in blood glucose regulation. Researchers have investigated neem extracts and individual constituents for possible effects on glucose metabolism, carbohydrate-digesting enzymes, oxidative stress and lipid-related parameters.

A critical review by Patil and colleagues examined experimental evidence concerning the antidiabetic potential of *Azadirachta indica*. The reviewed investigations included studies of blood glucose, glucose tolerance, lipid profiles and diabetic complications [4].

The available findings are promising for further investigation, but they do not establish neem as a replacement for standard antidiabetic treatment. Differences in experimental design, plant preparations and doses make direct comparisons difficult. Well-designed human studies are needed to determine clinical benefit, appropriate dosage, possible interactions with medicines and long-term safety.

6. Pharmaceutical and Industrial Applications:

Neem has potential applications across several fields because its plant parts provide chemically different materials. Its importance extends beyond traditional medicine to personal-care products, agricultural preparations and pharmaceutical research.

6.1 Herbal Formulations

Neem-derived ingredients have been incorporated into various traditional and commercial preparations. Depending on their intended use, these may include topical products and oral-care formulations. Product development requires appropriate selection of plant material, control of microbial contamination, stability testing and assessment of local tolerability.

6.2 Cosmetic and Personal-Care Products

Neem extracts and oils are used in selected skin-care, hair-care and oral-care products. Their use is supported by traditional practices and interest in their chemical properties. However, claims concerning the prevention or treatment of skin diseases should be supported by evidence appropriate to the specific finished product.

6.3 Agricultural Products

Neem-derived materials are important in the development of botanical insecticides and other agricultural products. Azadirachtin is particularly relevant because of its effects on insect feeding and development. Proper formulation and application are important to maintain effectiveness and reduce unintended environmental effects.

6.4 Pharmaceutical Research

The chemical diversity of neem provides opportunities for the isolation and characterization of compounds with potentially useful biological activities. Pharmaceutical research may focus on identifying active constituents, investigating their mechanisms, improving their stability and assessing whether they can be developed into safe and effective products.

7. Toxicity And Safety Considerations

The natural origin of a substance does not automatically make it safe. Neem preparations differ in chemical composition, concentration, route of administration and intended use. Consequently, the safety findings for one type of extract cannot necessarily be applied to all neem-derived products.

An updated review by Braga and colleagues examined the available in vivo toxicity evidence for *Azadirachta indica* and highlighted the importance of evaluating both mammalian and environmental toxicity [5]. The available evidence indicates that safety depends on the preparation, dose and exposure conditions.

Neem seed oil requires particular caution when considering oral exposure. Ingestion of neem oil has been associated with serious poisoning, especially in children. Therefore, neem oil should not be swallowed or used for self-treatment without appropriate medical guidance.

The use of neem during pregnancy also requires caution because adequate safety evidence is lacking for many preparations and routes of exposure. Products intended for medicinal use should undergo appropriate reproductive and developmental toxicity assessment when relevant.

Important safety considerations include:

- Correct identification and authentication of the plant material.
- Evaluation of contaminants, adulterants and microbial quality.
- Standardization of extraction and manufacturing procedures.
- Assessment of acute and repeated-dose toxicity.
- Investigation of possible interactions with conventional medicines.
- Evaluation of reproductive safety and risks in vulnerable populations.
- Appropriate labelling, storage and control of the intended route of administration.

Further toxicological and clinical studies are necessary to establish the safety limits of specific standardized preparations. Traditional use alone is insufficient to establish the safety of concentrated extracts or isolated compounds.

8. Conclusion

Azadirachta indica is an important medicinal tree with a long history of traditional use and a diverse range of phytochemical constituents. Its leaves, bark, seeds and other plant parts contain compounds that have attracted scientific interest for their possible antimicrobial, antioxidant, anti-inflammatory, antidiabetic and other biological activities. Neem is also important in agricultural pest management because of the insecticidal and antifeedant properties of azadirachtin and related constituents. The available experimental evidence supports continued investigation of neem as a source of potentially useful natural compounds. However, the biological activity of an extract depends on its composition, preparation and concentration. Findings from laboratory assays and animal studies should not be treated as proof of clinical efficacy, and the safety of one neem product should not automatically be extended to another. Further progress will depend on botanical authentication, chemical standardization, reproducible pharmacological testing, appropriate toxicological assessment and well-designed clinical studies. With these measures, research may help distinguish traditional claims from scientifically supported applications and identify neem-derived compounds suitable for future pharmaceutical development.

References

1. Cowan MM. Plant products as antimicrobial agents. Clin Microbiol Rev. 1999;12(4):564-582.
2. Draelos ZD. The science behind skin care: moisturizers. J Cosmet Dermatol. 2013;12(4):263-268.
3. Lodén M. Role of topical emollients and moisturizers in the treatment of dry skin barrier disorders. Am J Clin Dermatol. 2003;4(11):771-788.
4. Surjushe A, Vasani R, Saple DG. Aloe vera: a short review. Indian J Dermatol. 2008;53(4):163-166.
5. Hamman JH. Composition and applications of Aloe vera leaf gel. Molecules. 2008;13(8):1599-1616.
6. Biswas K, Chattopadhyay I, Banerjee RK, Bandyopadhyay U. Biological activities and medicinal properties of neem. Curr Sci. 2002;82(11):1336-1345.
7. Alzohairy MA. Therapeutics role of Azadirachta indica and their active constituents in diseases prevention and treatment. Evid Based Complement Alternat Med. 2016;2016:7382506.
8. Hewlings SJ, Kalman DS. Curcumin: a review of its effects on human health. Foods. 2017;6(10):92.
9. Agero ALCA, Verallo-Rowell VM. A randomized double-blind controlled trial comparing extra virgin coconut oil with mineral oil as a moisturizer. Dermatitis. 2004;15(3):109-116.
10. Intahphuak S, Khonsung P, Panthong A. Anti-inflammatory, analgesic, and antipyretic activities of virgin coconut oil. Pharm Biol. 2010;48(2):151-157.
11. Rawlings AV, Harding CR. Moisturization and skin barrier function. Dermatol Ther. 2004;17 Suppl 1:43-48.
12. Aulton ME, Taylor KMG, editors. Aulton's Pharmaceutics: The Design and Manufacture of Medicines. 6th ed. Elsevier; 2021.
13. Ansel HC, Allen LV, Popovich NG. Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems. 10th ed. Lippincott Williams & Wilkins; 2014.

14. Barel AO, Paye M, Maibach HI, editors. Handbook of Cosmetic Science and Technology. 4th ed. CRC Press; 2014.

15. Fluhr JW, Darlenski R, Surber C. Glycerol and the skin. Br J Dermatol. 2008;159(1):23-34.

16. Saleem S, Muhammad G, Hussain MA, Bukhari SNA. A comprehensive review of phytochemical profile, bioactives for pharmaceuticals, and pharmacological attributes of *Azadirachta indica*. Phytother Res. 2018;32(7):1241–1272. doi:10.1002/ptr.6076.

17. Gupta SC, Prasad S, Tyagi AK, Kunnumakkara AB, Aggarwal BB. Neem (*Azadirachta indica*): An Indian traditional panacea with modern molecular basis. Phytomedicine. 2017;34:14–20. doi:10.1016/j.phymed.2017.07.001.

18. Sarkar S, Singh RP, Bhattacharya G. Exploring the role of *Azadirachta indica* (neem) and its active compounds in the regulation of biological pathways: An update on molecular approach. 3 Biotech. 2021;11(4):178. doi:10.1007/s13205-021-02745-4.

19. Patil SM, Shirahatti PS, Ramu R. *Azadirachta indica* A. Juss (neem) against diabetes mellitus: A critical review on its phytochemistry, pharmacology, and toxicology. J Pharm Pharmacol. 2022;74(5):681–710. doi:10.1093/jpp/rgab098.

20. Braga TM, Rocha L, Chung TY, Oliveira RF, Pinho C, Oliveira AI, et al. *Azadirachta indica* A. Juss. In vivo toxicity—An updated review. Molecules. 2021;26(2):252. doi:10.3390/molecules26020252.

21. Fernandes SR, Barreiros L, Oliveira RF, Cruz C, Prudêncio C, Oliveira AI, et al. Chemistry, bioactivities, extraction and analysis of azadirachtin: State-of-the-art. Fitoterapia. 2019;134:141–150. doi:10.1016/j.fitote.2019.02.006.

22. Patel SM, Nagulapalli Venkata KC, Bhattacharyya P, Sethi G, Bishayee A. Potential of neem (*Azadirachta indica* L.) for prevention and treatment of oncologic diseases. Semin Cancer Biol. 2016;40–41:100–115. doi:10.1016/j.semcancer.2016.03.002.