



International Journal of Research Publication and Reviews

Journal Homepage: www.ijrpr.com ISSN 2582-7421

Readiness Determination in School-Based Trauma Intervention: How Clinicians Decide When a Child Can Safely Engage Trauma-Focused Treatment

Babatola Arogundade

University of Wisconsin-Oshkosh, USA

DOI : <https://doi.org/10.55248/gengpi.07.0926.3116>

ABSTRACT

Child trauma treatment endorses safety and stabilization before trauma processing, yet determining when a child is ready remains inadequately operationalized in school-based mental health care across the United States. Premature progression can precipitate symptom escalation, dissociation, shutdown, disengagement, or treatment dropout, while school schedules, group delivery, limited caregiver involvement, environmental instability, and variable trauma-specific preparation complicate clinical judgment. This paper examines readiness provisions across CBITS, Bounce Back, SSET, TF-CBT, SPARCS, ARC, TARGET, and EMDR, coding whether readiness criteria are explicit, implied, absent, or dependent upon clinician judgment. It distinguishes symptom measurement from readiness determination and identifies a related supervisory gap: verifying clinician capacity to make and document this decision. The paper introduces Creating Conditions for Becoming, a threshold-gated framework assessing affect regulation, environmental stability, supportive-adult availability, therapeutic alliance and willingness, dissociation, developmental capacity, and observable readiness loss. Progression requires documented supervisory verification rather than predetermined session counts. The framework provides a testable structure for clinical practice, supervision, and training, while establishing priorities for Delphi consensus, readiness-instrument validation, and school-based outcome research supporting safer trauma treatment nationally.

Keywords: Trauma Treatment Readiness; School-Based Mental Health; Creating Conditions for Becoming; Child Trauma; Clinical Supervision; Trauma-Focused Treatment

1. INTRODUCTION: THE UNRESOLVED READINESS DECISION

1.1 Readiness as the Clinical Pivot Point

A pivotal decision in child trauma treatment occurs before direct engagement with traumatic memories: whether a child is sufficiently prepared to tolerate trauma-focused processing without overwhelming existing regulatory capacities [1]. Readiness therefore marks the clinical boundary between stabilization and deliberate engagement with traumatic material. Although evidence-based interventions provide structured treatment sequences, progression through those sequences does not necessarily establish that an individual child possesses the emotional, relational, developmental, and environmental conditions required for processing [2]. Symptom severity is similarly insufficient. A child with substantial post-traumatic symptoms may retain meaningful regulatory capacity, whereas another with lower measured severity may respond to trauma activation with dissociation, shutdown, avoidance, or dysregulation [3].

Phase-oriented trauma theory provides an important foundation for this distinction. Herman's influential formulation places establishment of safety before remembrance and mourning and subsequent reconnection [4]. This sequencing has contributed to the enduring clinical principle that safety and stabilization should precede intensive engagement with traumatic experiences. Related trauma-treatment literature likewise emphasizes preparation, coping capacity, and therapeutic security before demanding trauma processing [3]. Yet agreement that clinicians should “establish safety first” does not resolve the operational question: what observable evidence demonstrates that sufficient readiness has actually been achieved? [5]. Without explicit thresholds, progression can remain dependent on clinician interpretation.

1.2 Consequences of Advancing Before Readiness

That ambiguity has clinical consequences. Trauma-focused work intentionally activates memories, emotions, cognitions, and physiological responses associated with traumatic experiences [4]. When activation exceeds a child's available regulatory capacity, the therapeutic process may produce escalating distress, affective flooding, avoidance, dissociation, behavioral shutdown, or disengagement [6]. Premature progression may also weaken

therapeutic trust or contribute to treatment interruption and dropout, particularly when children experience processing as something imposed rather than collaboratively undertaken [3].

SAMHSA's trauma-informed guidance warns against inadequately prepared practitioners pressing clients for trauma material without adequately understanding trauma responses and appropriate pacing [7]. The readiness problem therefore operates at two levels. Clinicians must determine whether the child is ready, but systems must also determine whether the clinician is competent to make that judgment [5]. Recognizing destabilization, differentiating tolerable therapeutic activation from loss of regulation, and knowing when to return to stabilization are clinical competencies rather than procedural assumptions [8].

1.3 Why Readiness Determination Matters Nationally in U.S. Schools

This distinction has particular national significance because schools constitute a major point of access to mental-health services for children and adolescents in the United States [1]. School-based services can reduce transportation, financial, stigma-related, and accessibility barriers that otherwise limit treatment utilization [6]. Their accessibility, however, does not eliminate the complexity of trauma treatment. School clinicians work within schedules, institutional demands, confidentiality requirements, variable caregiver participation, and environments where treatment pacing cannot always be determined solely by clinical need [8].

The resulting gap is specific. Trauma protocols increasingly describe what interventions clinicians should deliver, yet the threshold establishing when this particular child can safely progress may remain incompletely operationalized [4]. This article therefore compares readiness provisions across major child-trauma protocols, examines the distinction between symptom measurement and readiness determination, evaluates clinician preparation and supervisory verification, and introduces Creating Conditions for Becoming (CCFB) as a structured readiness framework [9]. CCFB reframes progression as a threshold-based clinical determination rather than an consequence of completed sessions.

If readiness is difficult to operationalize under conventional clinical conditions, school-based delivery introduces structural constraints that make individualized determination more difficult—and more necessary [7].

2. WHY SCHOOL-BASED TRAUMA TREATMENT COMPLICATES READINESS

2.1 Clinical Treatment Within an Educational Schedule

School-based trauma intervention improves accessibility by locating services where children spend substantial time, but it embeds clinical treatment within an institution organized around educational rather than therapeutic priorities [12]. Session duration and frequency may consequently be shaped by class periods, examinations, assemblies, attendance, holidays, transportation schedules, and competing school activities [15]. These constraints become clinically important when trauma interventions require sequential movement from psychoeducation and stabilization toward progressively greater engagement with traumatic material. Structured interventions such as CBITS demonstrate the advantages of predictable sequencing, yet scheduled progression and individual readiness are conceptually different decisions [10]. A child reaching the session designated for trauma processing has satisfied a temporal requirement, not necessarily a clinical threshold.

This distinction becomes particularly important following interruption. Missed sessions or shortened encounters can disrupt continuity, while institutional pressure to complete an intervention within an academic term may unintentionally compete with individualized pacing [17]. Readiness determination must therefore remain responsive to the child's current capacity rather than being inferred from protocol position or elapsed treatment time [13].

2.2 Environmental Safety, Caregiver Support, and Disclosure

Unlike residential or highly controlled treatment settings, schools cannot determine the conditions to which children return after each session. Trauma-exposed children may return to unstable caregiving arrangements, continuing adversity, community violence, family conflict, or other environments capable of undermining gains achieved during stabilization [11]. Safety must consequently be understood beyond the therapy room. Apparent regulation during a school session may coexist with environmental circumstances that substantially alter the child's capacity to tolerate subsequent trauma processing [16].

Caregiver involvement creates another challenge. Family participation can strengthen child trauma treatment through reinforcement of coping skills, improved communication, and support outside sessions [14]. Yet school-based programs frequently encounter practical barriers to sustained caregiver participation, making assumptions derived from clinic-based models difficult to reproduce consistently [18]. Confidentiality and mandated-reporting responsibilities can further influence disclosure. Children who anticipate consequences following disclosure may withhold information relevant to determining their current safety or readiness [12].

2.3 Group Delivery, Caseloads, and Unequal Readiness

Tier 2 group interventions create an additional tension between standardized delivery and individualized clinical need. Children participating in the same group may differ substantially in affect regulation, dissociation, developmental capacity, environmental stability, therapeutic trust, and willingness to discuss trauma [15]. Nevertheless, group protocols generally advance collectively, potentially creating a mismatch between group pacing and individual readiness [10].

School mental-health professionals may simultaneously manage substantial caseloads, crisis responsibilities, documentation demands, and students presenting with diverse clinical needs [17]. Moreover, practitioners delivering trauma-informed services are not invariably trauma specialists or recipients of continuous trauma-specific supervision [13]. Under these conditions, individualized readiness assessment can become difficult precisely where it is most necessary.

School-based treatment therefore introduces a fundamental clinical tension: the intervention may operate according to a schedule while readiness operates according to the child [16]. Resolving that tension requires examining whether existing trauma protocols provide observable criteria for progression, distinguish preparation from demonstrated readiness, and specify when processing should be delayed or stopped [18].

3. WHAT EXISTING TRAUMA PROTOCOLS ACTUALLY SAY ABOUT READINESS

3.1 Review Method and Readiness Coding Framework

Eight interventions were selected because they represent school-based, child-focused, complex-trauma, or processing approaches relevant to decisions about progression: CBITS, Bounce Back, SSET, TF-CBT, SPARCS, ARC, TARGET, and EMDR [12]. The review focuses on what each model requires before trauma material is directly processed, rather than comparing effectiveness. Protocol descriptions, manuals, implementation materials, and program documentation were examined for preparation activities, progression rules, contraindications, reassessment procedures, and stopping guidance [18]. Readiness was coded as specified when observable criteria governed progression; implied when stabilization or preparation preceded processing without an explicit threshold; absent when no identifiable readiness criterion was stated; and clinician judgment when progression depended primarily on individualized professional determination [14]. A second distinction classified progression as fixed-sequence when treatment advanced principally through predetermined components or sessions, or assessed-threshold when advancement depended upon demonstrated clinical conditions [21]. These categories permit preparation and readiness to be analyzed as related but non-equivalent constructs.

3.2 School-Based Structured Protocols: CBITS, Bounce Back, and SSET

CBITS illustrates the distinction clearly. Its school-based cognitive-behavioral structure combines psychoeducation, relaxation, cognitive restructuring, social problem-solving, and exposure to traumatic memories within an organized course of treatment [13]. Skills acquisition precedes or accompanies more direct trauma engagement, providing preparation before exposure. Nevertheless, the sequence itself does not constitute a standardized child-specific go/no-go instrument demonstrating that regulation, environmental stability, willingness, and tolerance have reached a defined threshold [17]. Accordingly, readiness is best characterized as implied within structured preparation, with important progression decisions remaining clinically mediated.

Bounce Back adapts this approach for younger elementary-school children. It combines ten group sessions emphasizing coping and practice with individual sessions used to process traumatic memories and grief [19]. Developmental adaptations and caregiver materials strengthen preparation, but the published structure again provides a treatment pathway rather than a universally scored readiness threshold [15].

SSET modifies CBITS for delivery by school personnel and reduces reliance on individual clinical and caregiver components [22]. Its accessibility is valuable in practice, yet this adaptation sharpens the readiness question because structured group progression may occur among children with unequal capacities for trauma engagement [12]. Across these three programs, preparation is evident; a uniform observable threshold that independently authorizes progression is less evident. The clinically important distinction is therefore between completing preparatory content and demonstrating readiness for the next therapeutic demand.

3.3 Clinical and Complex-Trauma Models: TF-CBT, SPARCS, ARC, and TARGET

TF-CBT provides a more explicitly phased architecture. Its PRACTICE components move through psychoeducation and parenting, relaxation, affect modulation, cognitive coping, trauma narration and processing, in-vivo mastery, conjoint sessions, and enhancement of safety [16]. Gradual exposure is integrated across treatment, and therapists calibrate exposure intensity according to what the child can tolerate without becoming overwhelmed [20]. This makes TF-CBT more responsive to individual tolerance than a simple session-count model. Even so, tolerance is clinically interpreted rather than established through a separate validated readiness score [14]. Guidance for complex trauma further recognizes fluctuating self-regulation and permits progression when youth have developed some usable coping capacity, illustrating both individualized pacing and the continuing importance of clinician judgment [18].

SPARCS approaches the issue differently. Designed for adolescents affected by chronic interpersonal trauma and ongoing stress, it is present-centered and emphasizes affect regulation, mindfulness, relationships, meaning, and coping rather than requiring detailed trauma narration as its organizing objective [13]. Its suitability for youth who may remain in stressful environments demonstrates why environmental safety cannot always be treated as a completed prerequisite [21].

ARC similarly organizes intervention around attachment, self-regulation, and developmental competency, making regulation and relational functioning central treatment targets [17]. As a flexible framework rather than a narrowly sequenced exposure protocol, it supports individualized formulation but does not provide a single standardized threshold authorizing trauma processing [15].

TARGET is likewise present-centered and teaches the FREEDOM sequence to strengthen emotional regulation, attentional control, executive functioning, mindfulness, and adaptive responses to stress [22]. Consequently, SPARCS, ARC, and TARGET offer substantial stabilization resources, yet their structures do not convert readiness into a common measurable gate. Across the four approaches, preparation and regulation are prominent; decisions concerning when particular trauma-processing demands are tolerable remain individualized and dependent upon case formulation, observed response, developmental needs, and professional judgment [16].

3.4 EMDR Preparation and Cross-Protocol Synthesis

EMDR makes preparation a distinct phase within its eight-phase model. Phase 2 emphasizes therapeutic alliance, informed understanding of the procedure, relaxation and affect-regulation resources, and sufficient stabilization to support subsequent reprocessing [20]. The clinician must therefore carefully evaluate whether the client can maintain adequate stability and dual attention as work moves toward assessment and desensitization [14].

Across all eight approaches, a consistent clinically important pattern therefore emerges. Preparation is rarely ignored: coping skills, regulation, psychoeducation, relational safety, resource development, or gradual exposure appear in different combinations [19]. What varies significantly is whether these elements function as treatment components, considerations, or explicit gates to progression. TF-CBT and EMDR articulate individualized tolerance more directly, whereas school-based protocols often embed preparation within structured sequences; SPARCS, ARC, and TARGET emphasize regulation and adaptive functioning without reducing readiness to a discrete score [13].

The resulting evidentiary distinction for readiness determination is fundamental: having a preparation phase is not equivalent to having an operational readiness criterion [21]. A protocol may teach regulation before processing while leaving unanswered how much regulation is sufficient or who verifies progression [15]. Thus, the cross-protocol gap is not an absence of stabilization practices. It is the absence of a shared, measurable, clinically defensible, clearly child-specific determination that separates “preparation completed” from “readiness demonstrated” [22]. This distinction provides the basis for examining whether existing assessment instruments can supply the missing determination.

Table 1. Protocol-Level Comparison of Readiness Determination in Child Trauma Interventions

Protocol	Population/setting	Stabilization/preparation component	Readiness characterization	Explicit threshold?	Progression structure	Primary decision-maker	Stop/ return criterion?
CBITS	School-aged youth; primarily school-based groups	Psychoeducation, relaxation, cognitive restructuring, coping and problem-solving before/directly alongside trauma exposure	Implied through preparatory skill development	No standardized child-specific threshold identified	Predominantly structured sequence with clinical adaptation	Clinician	Clinical response may require adaptation, but no uniform readiness-loss threshold
Bounce Back	Elementary-school children, approximately ages 5–11	Group coping-skills development with individual trauma-memory processing and caregiver-related components	Implied through preparation and developmental adaptation	No standardized readiness score identified	Structured group and individual sequence	Clinician	Individual clinical judgment
SSET	School-based youth; designed for delivery by school personnel	Psychoeducation, relaxation, cognitive coping, problem-solving and trauma-related activities	Implied	No explicit child-specific go/no-go threshold identified	Predominantly structured sequence	Facilitator/clinician within program structure	Limited explicit threshold for returning to stabilization

Protocol	Population/setting	Stabilization/preparation component	Readiness characterization	Explicit threshold?	Progression structure	Primary decision-maker	Stop/ return criterion?
TF-CBT	Trauma-exposed children/adolescents with caregiver involvement	PRACTICE stabilization components; relaxation, affect modulation, cognitive coping and gradual exposure	Implied with substantial clinician judgment; tolerance is actively evaluated	Individual tolerance is considered, but not reduced to a standardized readiness score	Phased and clinically individualized	Trained clinician with child/caregiver participation	Pacing and exposure intensity modified according to tolerance
SPARCS	Adolescents with chronic/complex trauma, including continuing stress	Present-centered affect regulation, mindfulness, interpersonal and coping skills	Implied/contextual	No discrete processing-readiness threshold identified	Structured skills-based sequence with individualized application	Clinician/group facilitator	Regulation and current functioning inform clinical response
ARC	Children/adolescents with complex trauma across service settings	Attachment, regulation and competency domains	Clinician judgment within framework	No universal processing threshold identified	Flexible framework rather than fixed exposure sequence	Clinician/team	Intervention may return to regulatory or relational targets as clinically indicated
TARGET	Adolescents/adults; individual, group and systems applications	FREEDOM skills addressing regulation, attention, executive functioning and adaptive stress responses	Implied through regulatory capacity	No standardized trauma-processing readiness threshold identified	Manualized skills sequence	Clinician/facilitator	Response guided by regulation and functioning
EMDR	Children, adolescents and adults; clinical settings	Phase 2 preparation, alliance, psychoeducation, stabilization, affect regulation and resource development	More explicitly readiness-oriented but clinician determined	Readiness is addressed explicitly, but determination remains individualized rather than a universal scored threshold	Phase-based progression informed by clinical assessment	EMDR-trained clinician with client	Additional preparation/stabilization when adequate readiness is not established

The protocol comparison demonstrates that preparation and stabilization are prominent across trauma interventions, but their presence does not automatically produce a common, measurable readiness threshold. The next question is therefore whether instruments routinely used in child trauma assessment can fill that gap. If those measures quantify symptoms rather than capacity to tolerate processing, the measurement problem becomes central to readiness determination.

4. THE MEASUREMENT PROBLEM: SYMPTOMS ARE NOT READINESS

4.1 What Existing Child Trauma Measures Actually Measure

Assessment instruments used in child trauma practice provide essential information, but their intended constructs must be distinguished from treatment readiness. The UCLA PTSD Reaction Index assesses trauma exposure and post-traumatic stress symptoms in children and adolescents, including symptom frequency and associated impairment [21]. The Child PTSD Symptom Scale similarly measures post-traumatic stress symptom severity and can support screening and monitoring of treatment-related change [19]. The Trauma Symptom Checklist for Children extends assessment beyond PTSD to domains including anxiety, depression, anger, post-traumatic stress, dissociation, and sexual concerns [24]. The Strengths and Difficulties Questionnaire provides a broader behavioral and emotional screening profile encompassing emotional symptoms, conduct difficulties, hyperactivity,

peer relationships, and prosocial behavior [22]. These instruments serve clinically valuable purposes and should not be evaluated against functions for which they were not designed [25]. Collectively, they identify symptoms, difficulties, and changes in functioning; they do not, by themselves, establish whether trauma processing should begin.

4.2 Symptom Severity Versus Capacity to Engage

The distinction is conceptual as well as practical: symptom severity asks how much distress or impairment is present; readiness asks whether this particular child, under current conditions, can safely tolerate the particular processing task being proposed [20]. A high PTSD score may indicate substantial clinical need without demonstrating that the child lacks sufficient capacity for carefully paced processing [23]. Conversely, a comparatively low symptom score cannot establish that the child possesses adequate affect regulation, environmental stability, therapeutic trust, or developmental capacity for trauma-focused engagement [19].

Dissociation illustrates the problem particularly clearly. Although dissociative symptoms can be measured within broader trauma assessment, their clinical significance during trauma processing concerns whether the child can remain sufficiently present and regulated while traumatic material is activated [24]. Similar distinctions apply to willingness, alliance, caregiver or supportive-adult availability, and current living conditions [21]. These variables concern the circumstances under which treatment occurs, not simply symptom magnitude. Readiness must therefore be conceptualized as a multidimensional and potentially fluctuating clinical state rather than as an extension of symptom severity [25].

4.3 The Missing Readiness Construct

Within the measures examined, no instrument provides an integrated determination combining regulatory capacity, environmental stability, relational support, willingness, dissociation, and developmental capacity into a specific decision about progression to trauma processing [22]. This finding should be interpreted narrowly. It does not establish that no relevant pediatric readiness measure exists anywhere in the literature, nor does it diminish validated instruments designed for screening, diagnosis, symptom monitoring, or functional assessment [20]. Rather, the reviewed measurement landscape reveals a construct-level gap between assessing trauma-related clinical presentation and determining readiness for a particular trauma-processing task [23].

That distinction identifies an empirical question requiring systematic investigation: whether readiness can be operationalized as a sufficiently coherent construct to support reliable clinical determination [25]. Until such evidence is established, symptom scores should inform readiness judgments without being treated as substitutes for them [21].

Figure 1. Symptom Severity and Trauma-Treatment Readiness as Distinct Clinical Constructs




Figure 1. Symptom Severity and Trauma-Treatment Readiness as Distinct Clinical Constructs

If standardized symptom instruments describe clinical presentation without independently determining readiness, responsibility for deciding whether processing should proceed returns substantially to professional judgment. The adequacy of clinician preparation and the capacity of supervision to examine that judgment therefore become central components of the safety question.

5. WHO MAKES THE READINESS DECISION? CLINICAL COMPETENCY AND SUPERVISION

5.1 *Readiness as an Unstandardized Clinical Judgment*

When protocols and symptom measures do not provide a definitive progression threshold, readiness ultimately depends substantially on clinical judgment [24]. Counselor preparation literature has long recognized difficulty defining minimum competence because competence encompasses knowledge, skills, professional judgment, ethical conduct, and the capacity to respond appropriately to individual clients [27]. General counseling competence, however, should not be conflated with trauma-treatment competence. Trauma-focused practice additionally requires knowledge of traumatic stress, stabilization, dissociation, developmental responses, pacing, and risks associated with activating traumatic material [23]. Readiness determination represents a still narrower competency: the ability to decide whether a particular child, under current conditions, can tolerate a proposed processing task and what evidence justifies that conclusion [29]. A clinician may therefore be generally competent and trauma-informed without having demonstrated a reproducible method for making this specific decision. Where shared operational criteria are limited, professional judgment remains indispensable, but its basis may remain difficult for another practitioner to examine [25].

5.2 *Supervision as the Clinical Checkpoint*

Clinical supervision is the most immediate professional setting in which readiness judgments can be examined before consequential treatment decisions are implemented [28]. Effective trauma supervision can support case conceptualization, ethical decision-making, clinician development, recognition of countertransference, and management of emotionally demanding clinical work [26]. Yet supervision availability does not necessarily mean that readiness determination itself is systematically verified. A supervisor may discuss symptoms, risk, treatment progress, and clinician reactions without establishing whether the clinician can assess each condition required for trauma processing [30].

A readiness-focused checkpoint would ask whether the clinician can distinguish symptom severity from readiness, identify dissociation or destabilization, explain why progression is justified, recognize observable evidence that readiness has been lost, and specify the response required when that occurs [24]. It would also test whether the clinician can delay progression when a school calendar, caseload expectation, caregiver preference, or protocol timetable favors advancement [27]. Supervision would thereby examine the reasoning supporting progression.

5.3 *The Verification Gap*

Existing trauma-supervision competencies address important but distinguishable concerns. NCTSN guidance concerning secondary traumatic stress, for example, emphasizes organizational and supervisory responses that help professionals recognize and manage the effects of indirect trauma exposure arising from their work [25]. Such competencies protect workforce functioning and quality of care, but they do not inherently establish that a supervisor has verified a clinician's competence to determine when a child can safely begin trauma processing [30].

The distinction is consequential. Supporting a clinician who is affected by trauma work asks whether the practitioner can remain effective and adequately supported; readiness verification asks whether that practitioner can make and defend a child-specific progression decision [23]. The latter requires observable evidence of assessment, rationale, pacing, contraindication recognition, and response to lost readiness [28]. This article defines the absence of a consistent mechanism for confirming those capacities as the readiness-verification gap [26]. Closing that gap requires moving supervision from assumed competence toward documented clinical verification.

The significance of the readiness-verification gap becomes clearer when treatment attrition and premature termination are considered. Existing outcome systems may document what happened during treatment without establishing whether readiness determination or clinician preparedness contributed to that trajectory.

6. EVIDENCE OF CONSEQUENCE AND THE MISSING LINK IN OUTCOME DATA

6.1 *Attrition and Premature Termination*

Treatment attrition is well documented in child and adolescent mental-health services, including services for trauma-exposed youth, but its causes are heterogeneous [27]. The National Child Traumatic Stress Network Core Data Set created an unusually large clinical record describing trauma histories, symptoms, functional difficulties, services, and outcomes among children receiving trauma-related care across diverse United States settings [23]. Analyses of these data demonstrate substantial clinical complexity, including multiple trauma exposures, behavioral difficulties, post-traumatic symptoms, and variation in service trajectories [29]. Other trauma-treatment studies likewise identify premature termination as a continuing implementation concern, with engagement influenced by child, caregiver, treatment, provider, and contextual factors [25].

These findings should not be interpreted as evidence that inadequate readiness determination causes dropout. Such a causal conclusion would require readiness to have been independently defined, measured, and prospectively related to termination [30]. The more precise question is whether existing datasets measured readiness sufficiently to test that hypothesis. Where readiness is not recorded as a distinct construct, its contribution to treatment continuation, destabilization, or premature termination cannot be isolated retrospectively [24].

6.2 The Readiness–Outcome Inversion

This limitation produces what may be described as a readiness–outcome inversion. Large clinical datasets can document the child's trauma history, symptoms, services received, treatment trajectory, and subsequent outcomes [28]. Conceptually, the recorded sequence can therefore encompass symptoms, treatment, completion or termination, and outcomes. This architecture is valuable for understanding service utilization and clinical change, but it may leave an earlier decision largely invisible [26].

The missing sequence concerns clinician readiness competence, child readiness determination, rationale for progression, observable loss-of-readiness signals, and clinician response [30]. Unless these variables are prospectively documented, researchers cannot readily determine whether a child entered trauma processing after demonstrated readiness, whether readiness changed during treatment, or how the clinician responded when tolerance deteriorated [23].

This inversion has methodological consequences. Poor outcomes cannot legitimately be attributed to readiness decisions when those decisions were never measured; equally, favorable outcomes cannot establish that readiness was assessed systematically [29]. The empirical problem is therefore one of missing explanatory information. A readiness framework must make the decision observable before research can determine whether its quality predicts clinically meaningful trajectories [25].

Table 2. The Readiness Evidence Gap Across Assessment, Treatment, Supervision, and Outcomes

Decision domain	What is currently documented	What is not consistently documented	Clinical consequence	Proposed CCFB response
Assessment	Trauma exposure, PTSD symptoms, emotional/behavioral difficulties, impairment and risk	Integrated determination of whether the child can tolerate the proposed trauma-processing task	Symptom severity may be mistaken for evidence for or against progression	Assess readiness independently across defined clinical conditions
Treatment preparation	Psychoeducation, coping skills, affect regulation, relaxation and stabilization activities	Observable threshold establishing when preparation has become sufficient readiness	Completion of preparatory components may be treated as equivalent to readiness	Require threshold-based progression rather than completion-based progression
Progression decision	Treatment phase, intervention delivered and sometimes clinician formulation	Explicit rationale documenting why the individual child was considered ready at that time	Basis for beginning processing may remain implicit	Record criteria, evidence and rationale before progression
Supervision	Case discussion, clinician support, risk management and treatment planning	Verification that the clinician can competently assess and defend readiness	Supervisor involvement does not necessarily demonstrate readiness competence	Require supervisor verification and countersigned readiness documentation
During processing	Symptoms, clinical observations and treatment response	Standardized loss-of-readiness indicators and predetermined responses	Destabilization may be recognized inconsistently or after escalation	Specify observable stop signals and required return-to-stabilization responses
Outcome	Completion, termination, symptom change and functional outcomes	Link between readiness determination, progression decisions and subsequent trajectory	Readiness cannot readily be examined as an explanatory clinical variable	Preserve longitudinal readiness records for outcome analysis

7. PROPOSED DIMENSIONS OF TRAUMA-TREATMENT READINESS

7.1 Regulation, Dissociation, and Developmental Capacity

Readiness should first incorporate behaviorally demonstrated affect-regulation capacity, rather than relying exclusively on a child's report of feeling prepared. Observable indicators may include the ability to recognize escalating distress, use previously learned regulation strategies, recover following

activation, and remain sufficiently engaged with the clinician [31]. Regulation does not require absence of distress; trauma processing inherently involves emotional activation [28]. The relevant question is whether activation remains within a range that permits therapeutic participation and recovery [34]. Dissociation requires separate consideration because apparent quietness or reduced emotional expression may reflect disengagement rather than regulation [29]. Clinicians must therefore distinguish regulated engagement from detachment, altered awareness, or diminished responsiveness. Developmental capacity is equally task-specific [33]. Readiness for verbal narration, cognitive restructuring, imagery, or other processing procedures depends partly on the child's cognitive, linguistic, emotional, and reflective abilities. Readiness consequently concerns demonstrated capacity for the particular therapeutic demand proposed [27].

7.2 Environmental and Relational Conditions

Readiness cannot be located entirely within the child because trauma processing occurs within relational and environmental systems [30]. Stability of the child's current living environment or placement should therefore be considered when determining whether gains achieved during sessions can be sustained outside treatment [35]. Instability does not automatically prohibit trauma-focused work, particularly when adversity cannot be rapidly changed, but it should influence formulation, pacing, safety planning, and monitoring [32]. The availability of at least one dependable supportive adult may similarly strengthen continuity between therapeutic sessions [28]. Within treatment, therapeutic alliance provides the relational context in which difficult material can be approached without the child experiencing processing as coercive [31]. The child's expressed willingness must therefore carry substantive weight. Agreement obtained through compliance, adult pressure, or perceived obligation is clinically different from collaborative willingness to participate [33]. Readiness is consequently both individual and contextual.

7.3 Readiness as Dynamic Rather Than Permanent

Readiness should also be understood as dynamic rather than permanently achieved. A child who demonstrates sufficient capacity during one session may return with reduced tolerance following renewed trauma exposure, family disruption, placement change, interpersonal conflict, sleep disturbance, or other destabilizing experiences [35]. Clinicians therefore require observable indicators of loss of readiness, including sustained dysregulation, marked dissociation, inability to remain oriented, escalating avoidance, withdrawal of willingness, or failure to recover following therapeutic activation [29]. Such signals should initiate predefined responses rather than improvised decisions [34]. Depending on presentation, the clinician may pause processing, reassess current conditions, restore stabilization work, modify pacing, consult supervision, strengthen external supports, or return temporarily to an earlier treatment phase [27]. Reassessment protects against treating a previous readiness determination as continuing authorization for processing [32]. Readiness is better conceptualized as a condition that must be established, monitored, and re-established when clinically necessary [30].

Transition. These dimensions become clinically useful only when organized into a reproducible decision pathway governing both progression and stopping. This requirement provides the conceptual basis for Creating Conditions for Becoming, which translates readiness from a collection of considerations into a threshold-gated clinical process.

8. CREATING CONDITIONS FOR BECOMING: A READINESS-GATED FRAMEWORK

8.1 Five-Stage Clinical Sequence

Creating Conditions for Becoming (CCFB) is proposed as a readiness-gated framework that organizes the preceding dimensions into five clinically linked stages: Conditions Assessment, Establishing Conditions, Verified Readiness, Graduated Engagement With Trauma Material, and Consolidation. Conditions Assessment identifies regulatory, dissociative, developmental, environmental, relational, and willingness factors relevant to the individual child [31]. Establishing Conditions addresses identified deficiencies before processing progresses [28]. Verified Readiness requires affirmative evidence that the applicable conditions have reached a clinically defensible threshold [35]. Graduated Engagement introduces trauma material incrementally while continuing readiness monitoring, rather than treating authorization to begin as permanent [29]. Consolidation evaluates recovery, integration, and continuing stability. Each stage contains a stop criterion: when required conditions are not demonstrated or subsequently deteriorate, progression pauses and the clinician stabilizes, modifies, or reassesses before proceeding [33].

8.2 Supervisor Verification and the Dated Readiness Record

CCFB additionally requires supervisory verification before direct trauma processing begins. The supervisor confirms that the clinician can assess each readiness dimension, distinguish symptom severity from readiness, recognize clinically significant dissociation, and identify observable loss-of-readiness signals [30]. The clinician must also articulate the required response to those signals and provide a defensible rationale for treatment pacing [34]. Importantly, competence includes the ability to delay progression despite pressure arising from caseload demands, caregiver expectations, protocol schedules, or school calendars [27]. The determination is entered as a dated, criteria-based readiness record countersigned in supervision, converting supervisory discussion into an examinable account of what was assessed, concluded, and authorized [32].

8.3 Accountability Across Clinical Transitions

Documentation becomes particularly important when care crosses organizational or professional boundaries. Children may change clinicians, supervisors, schools, placements, agencies, or service systems while trauma treatment remains incomplete [35]. Without an accessible record, a subsequent practitioner may know which intervention was delivered without knowing why processing began or whether readiness later deteriorated [28]. CCFB preserves the clinical reasoning behind progression, identified vulnerabilities, stop signals, and previous responses [31]. Readiness thereby becomes a transferable clinical determination rather than an undocumented impression that disappears when personnel or settings change [33].

Figure 2. Creating Conditions for Becoming: Threshold-Gated Readiness Determination Framework

9. IMPLICATIONS FOR U.S. PRACTICE, SUPERVISION, TRAINING, AND POLICY

9.1 Clinical Practice and Supervision

For school-based clinicians, CCFB could provide a feasible structure for converting an otherwise implicit pacing decision into an explicit clinical determination [35]. Rather than requiring readiness to be inferred from completed sessions, clinicians would examine defined conditions before progression and reconsider them when circumstances change [38]. The framework could simultaneously provide supervisors with shared terminology for questioning why processing should begin, continue, pause, or return to stabilization [34]. Such structure does not eliminate professional judgment; it makes the evidence supporting that judgment visible and reviewable [40]. In school environments where time and caseload pressures can influence pacing, this transparency could strengthen consistent clinical decision-making [36].

9.2 Counselor Preparation and Trauma-Specific Training

Independent decisions about trauma processing should follow preparation extending beyond general counseling competence [37]. Training should develop the clinician's capacity to assess readiness, recognize dissociation, evaluate environmental safety, adapt processing tasks developmentally, and distinguish tolerable activation from destabilization [39]. Clinicians should also demonstrate competence in pacing, documenting progression decisions, identifying loss-of-readiness signals, and specifying when treatment should return to stabilization [35]. Supervised practice could require trainees to articulate the clinical evidence supporting both progression and non-progression decisions [40]. This approach treats readiness determination as a demonstrable trauma-specific competency rather than an assumed consequence of professional qualification, protocol familiarity, or completion of general trauma-informed training [34].

9.3 United States Policy and Trauma-System Quality

At the systems level, CCFB aligns conceptually with the United States commitment to strengthening trauma services for children and families [36]. The Children's Health Act of 2000 authorized development of the National Child Traumatic Stress Initiative, supporting improved treatment, service development, training, and dissemination of evidence-based trauma knowledge [38]. Within that national objective, operationalizing readiness could add a quality dimension concerning when and under what conditions trauma processing proceeds [37]. If subsequently validated, readiness criteria and supervisory verification could support safer trauma engagement, stronger stabilization, improved treatment retention, more interpretable symptom trajectories, and treatment completion [39]. These remain proposed clinical benefits requiring empirical evaluation, not established effects of CCFB [35].

10. LIMITATIONS AND RESEARCH AGENDA

10.1 Limits of the Protocol Review

The protocol comparison identifies how selected interventions address preparation, stabilization, progression, and clinical judgment, but it cannot independently establish what readiness criteria should govern trauma processing [36]. Differences among protocols also reflect distinct populations, theoretical models, settings, and therapeutic objectives [39]. CCFB should therefore be understood as a proposed conceptual framework rather than a validated clinical standard [34]. Its dimensions and thresholds require systematic evaluation before they can support prescriptive decisions in routine child-trauma practice [38].

10.2 Delphi Consensus and Instrument Development

A logical next step is a Delphi study involving child-trauma clinicians, play-therapy supervisors, school-based practitioners, and trauma-treatment supervisors [40]. Successive consensus rounds could evaluate proposed readiness dimensions, operational thresholds, contraindications, loss-of-readiness indicators, and required clinical responses [35]. Findings could inform development of a Child Trauma Treatment Readiness Determination Instrument and companion Supervisor Readiness Verification Protocol [37]. Psychometric evaluation should examine internal consistency, interrater

reliability across clinicians and supervisors, and construct and discriminant validity [39]. Comparisons with established child-trauma measures would test whether readiness behaves as a distinguishable construct rather than simply reproducing PTSD severity, general distress, dissociation, or functional impairment [34].

10.3 Clinical Testing of Threshold-Gated Progression

Following consensus and preliminary validation, CCFB should be tested prospectively in United States school-based settings [38]. Single-case experimental designs with repeated between-session measurement could examine whether threshold-gated progression produces different clinical trajectories from fixed or conventionally sequenced stabilization [36]. Outcomes should remain clinically focused, including symptom trajectory, dysregulation, dissociation, treatment interruption, attrition, stabilization, and completion [40]. Repeated readiness measurement would additionally permit investigation of whether changes in readiness precede meaningful changes in treatment tolerance, providing an empirical test of the framework's proposed clinical mechanism [35].

Figure 3. Research and Validation Pathway for Creating Conditions for Becoming




Figure 3. Research and Validation Pathway for Creating Conditions for Becoming

11. CONCLUSION

The unresolved question in child trauma treatment is not whether stabilization should precede trauma processing, but how a clinician determines and demonstrates that sufficient readiness exists for a particular child at a particular time. This problem becomes especially consequential in school settings, where institutional schedules, environmental uncertainty, variable caregiver involvement, group delivery, and competing demands can complicate individualized clinical pacing.

The analysis demonstrates an important distinction between providing stabilization and establishing readiness. Existing trauma protocols offer substantial preparation strategies, yet readiness is not consistently operationalized as an observable threshold. Likewise, symptom measures characterize distress and impairment but do not independently establish capacity for trauma processing. Consequently, considerable responsibility remains with clinician judgment and supervision, exposing a readiness-verification gap.

Creating Conditions for Becoming addresses this gap by organizing readiness around regulatory, developmental, environmental, relational, and dissociative conditions and requiring threshold-based progression, reassessment, and supervisory documentation. CCFB does not replace CBITS, Bounce Back, SSET, TF-CBT, SPARCS, ARC, TARGET, or EMDR; it provides a decision and verification structure surrounding progression into trauma-focused work. Following systematic consensus, validation, and clinical testing, such a framework could strengthen clinical consistency, supervisory accountability, treatment safety, and quality of trauma care for children across United States schools.

REFERENCE

1. Wood CE, Domingues J, Howard JM, Said S. Understanding and responding to crisis and trauma in the school setting. In *Handbook of Australian school psychology: Integrating international research, practice, and policy* 2017 Jan 28 (pp. 413-434). Cham: Springer International Publishing.
2. Pittman JC, Bekele Y, Scardamalia K, Hoover S, Schaeffer CM. Addressing student trauma and loss: An evaluation of two school-based treatment models. *Children & Schools*. 2025 Jul;47(3):191-9.
3. Meldrum M, Gouze KR, Russo J, Kerns C, Holley C, Centola HJ, Sroka A, Cutshaw O, Cicchetti C. Meeting the Relational and Developmental Needs of Young Children Exposed to Potentially Traumatic Events Through Trauma-Informed Care. *Early Education and Development*. 2025 May 19;36(4):767-92.
4. Hodas GR. Responding to childhood trauma: The promise and practice of trauma informed care. Pennsylvania Office of Mental Health and Substance Abuse Services. 2006 Feb;177:5-68.
5. Sun Y, Blewitt C, Minson V, Bajayo R, Cameron L, Skouteris H. Trauma-informed interventions in early childhood education and care settings: A scoping review. *Trauma, Violence, & Abuse*. 2024 Jan;25(1):648-62.
6. Padden JV. Urban trauma: Mental health clinicians' perception of trauma informed care, trauma responsive practice, and trauma resilience. Morgan State University; 2021.
7. Morton E. Trauma-informed school practices: Building expertise to transform schools. George Fox University Library; 2019.
8. Davis W, Petrovic L, Whalen K, Danna L, Zeigler K, Brewton A, Joseph M, Baker CN, Overstreet S, New Orleans Trauma-Informed Learning Collaborative. Centering trauma-informed approaches in schools within a social justice framework. *Psychology in the Schools*. 2022 Dec;59(12):2453-70.
9. Baweja S, Santiago CD, Vona P, Pears G, Langley A, Kataoka S. Improving implementation of a school-based program for traumatized students: Identifying factors that promote teacher support and collaboration. *School Mental Health*. 2016 Mar;8(1):120-31.
10. Williams AM. *Trauma and Resiliency Program: A Program Design for Trauma-Exposed Elementary School-Aged Children in Rural Areas* (Doctoral dissertation, Capella University).
11. Mancini MA. A Pilot Study Evaluating a School-Based, Trauma-Focused Intervention for Immigrant and Refugee Youth: MA Mancini. *Child and Adolescent Social Work Journal*. 2020 Jun;37(3):287-300.
12. Plumb JL, Bush KA, Kerseviz SE. Trauma-sensitive schools: An evidence-based approach. *School Social Work Journal*. 2016 Mar 1;40(2):37-60.
13. Pressimone Y. *Trauma in Schools: Assessing Teacher Preparedness and Attitudes About Trauma-Informed Care* (Doctoral dissertation, Georgian Court University).
14. Kwaku Gyamfi Boamah. AI-Driven Threat Intelligence Adoption in Security Operations Centers: Human Skill Erosion or Analyst Augmentation? A Comparative Case Study of Resource-Constrained SOC's in Emerging Markets. *International Journal of Science and Research Archive*, 2026, 19(03), 025-033. Article DOI: <https://doi.org/10.30574/ijrsra.2026.19.3.1226>.
15. Connors EH, Prout J, Vivrette R, Padden J, Lever N. Trauma-focused cognitive behavioral therapy in 13 urban public schools: Mixed methods results of barriers, facilitators, and implementation outcomes. *School Mental Health*. 2021 Dec;13(4):772-90.
16. Ritblatt SN, Hokoda A, editors. From trauma to resiliency: Trauma-informed practices for working with children, families, schools, and communities. Taylor & Francis; 2022 Sep 28.
17. Chafouleas SM, Koriakin TA, Roundfield KD, Overstreet S. Addressing childhood trauma in school settings: A framework for evidence-based practice. *School mental health*. 2019 Mar 15;11(1):40-53.
18. Herrenkohl TI, Hong S, Verbrugge B. Trauma-informed programs based in schools: Linking concepts to practices and assessing the evidence. *American Journal of Community Psychology*. 2019 Dec;64(3-4):373-88.

19. Rishel CW, Tabone JK, Hartnett HP, Szafran KF. Trauma-informed elementary schools: Evaluation of school-based early intervention for young children. *Children & Schools*. 2019 Oct 10;41(4):239-48.
20. Wong M, Villaverde V, Vona P. Multi-tiered trauma focused interventions for safer schools. *The school services sourcebook: A guide for school-based professionals*. 2024:389-401.
21. Distel LM, Torres SA, Ros AM, Brewer SK, Raviv T, Coyne C, Baker S, Kolski C, Smith ML, Santiago CD. Evaluating the implementation of Bounce Back: Clinicians' perspectives on a school-based trauma intervention. *Evidence-Based Practice in Child and Adolescent Mental Health*. 2019 Jan 2;4(1):72-88.
22. Nadeem E, Saldana L, Chapman J, Schaper H. A mixed methods study of the stages of implementation for an evidence-based trauma intervention in schools. *Behavior therapy*. 2018 Jul 1;49(4):509-24.
23. Martin SL, Ashley OS, White L, Axelson S, Clark M, Burrus B. Incorporating trauma-informed care into school-based programs. *Journal of school health*. 2017 Dec;87(12):958-67.
24. Powell BJ, Patel SV, Haley AD, Haines ER, Knocke KE, Chandler S, Katz CC, Seifert HP, Ake III G, Amaya-Jackson L, Aarons GA. Determinants of Implementing Evidence-Based Trauma-Focused Interventions for Children and Youth: A Systematic Review: BJ Powell et al. *Administration and Policy in Mental Health and Mental Health Services Research*. 2020 Sep;47(5):705-19.
25. Hoffman RD. *Clinician perspectives on implementing a school-based trauma intervention for elementary school students: A mixed methods study of adaptations* (Doctoral dissertation, Rutgers University-Graduate School of Applied and Professional Psychology).
26. Delgaty LE. *Learning to Heal: Working Toward Trauma-Informed Frameworks in Hospital-Based Schools* (Doctoral dissertation, The University of North Carolina at Chapel Hill).
27. Alebiosu O. AI and cybersecurity governance and legal compliance challenges affecting digital energy infrastructure, smart grids, and critical utility resilience systems. *Int J Comput Appl Technol Res*. 2026;15(8):1-14. doi:10.7753/IJCATR1508.1001.
28. Hanrahan AR. *School Psychologists' Beliefs, Perceived Competence, Knowledge, Social/Norms Expectations, and Implementation of School-based Trauma-Focused Interventions* (Doctoral dissertation, The University of Wisconsin-Milwaukee).
29. Luberenga I. Addressing Childhood Trauma Through School-Based Mental Health Interventions: A Scoping Review of Trauma-Informed Approaches and Psychosocial Outcomes Among Vulnerable Youth in United States of America (USA).
30. Avery F, Kennedy N, James M, Jones H, Amos R, Bellis M, Hughes K, Brophy S. A systematic review of non-clinician trauma-based interventions for school-age youth. *PloS one*. 2024 Sep 6;19(9):e0293248.
31. McNerny K. *Understanding Successful Trauma-Informed Implementation in Schools: A Study of School Contexts and Needs* (Doctoral dissertation, Rutgers The State University of New Jersey, Graduate School of Applied and Professional Psychology).
32. Abdullahi Bamidele Bashiru; The Social Resource Network (SRN): A Socio-Technical Framework for Circular Waste Matchmaking. *ACS Sustainable Resour. Manage*. 26 February 2026; 3 (2): 283–285. <https://doi.org/10.1021/acssusresmgt.5c00510>
33. Dombo EA, Sabatino CA. Creating trauma-informed schools: A guide for school social workers and educators. Oxford University Press; 2019 Jan 15.
34. Nadeem E, Ringle V. Culturally Responsive, Trauma-Informed Care for Newcomer Students: Engagement and Clinical Strategies from Providers in School-Based Health Centers. *Journal of Trauma Studies in Education*. 2025;4(3):151-79.
35. Bekele Y, Scardamalia K, Hoover SA, Schaeffer CM. Who Gets Referred to School-Based Trauma Treatment? A Descriptive Analysis of Bounce Back and Support for Students Exposed to Trauma. *Psychology in the Schools*. 2025 Oct;62(10):4195-206.
36. Blodgett C, Dorado J. A selected review of trauma-informed school practice and alignment with educational practice. CLEAR Trauma Center. San Francisco, CA: University of California. 2016;1:1-88.
37. Jaycox LH, Kataoka SH, Stein BD, Wong M, Langley A. Responding to the needs of the community: A stepped care approach to implementing trauma-focused interventions in schools. *Report on Emotional and Behavioral Disorders in Youth*. 2005;5(4):85-8.
38. Chafouleas SM, Johnson AH, Overstreet S, Santos NM. Toward a blueprint for trauma-informed service delivery in schools. *School mental health*. 2016 Mar;8(1):144-62.
39. Fitzgerald MM, Cohen JA. Trauma-focused cognitive behavior therapy for school psychologists. *Journal of Applied School Psychology*. 2012 Jul 1;28(3):294-315.
40. Lopez A. *Evidence-Based Practice or Practice-Based Evidence? A Qualitative Examination of School Psychologists' Implementation of Trauma-Informed Interventions* (Doctoral dissertation).